

Detection of CRABP1 in Formalin-Fixed, Paraffin-Embedded Mouse Tissue

Reagent and Antibody Information

[1X Wash Buffer](#)

[3% Hydrogen Peroxide](#)

[1% BSA Diluent](#)

[1X EDTA](#)

[DAB Chromogen](#)

[Hematoxylin](#)

Blocking Solution: Rodent Block M

Biocare Medical
Concord, CA 94520
www.biocare.net
1-800-799-9499
Catalog # RBM961

Primary Antibody: Mouse Monoclonal Antibody To CRABP1

Abcam, Inc
Cambridge, MA 02139
www.abcam.com
1-888-772-2226
Catalog # ab2816

Negative Control Serum: Purified Mouse IgG2b Isotype Control Serum

BD Biosciences
San Jose, CA 95131
www.bdbiosciences.com
1-855-236-2772
Catalog # 557351

Polymer Reagent: Mouse-on-Mouse HRP-Polymer Detection

Biocare Medical
Concord, CA 94520
www.biocare.net
1-800-799-9499
Catalog # MM620

Staining Procedure

Positive Control Tissue: Embryo

Stain Localization: Cytoplasmic / nuclear

1. Deparaffinize and hydrate slides through the following solutions:

Solution	Repetitions	Time
Xylene	2 times	5 minutes
100% Ethanol	2 times	3 minutes
95% Ethanol	2 times	3 minutes
1X Wash Buffer	2 times	5 minutes

2. Quench endogenous peroxidase by placing the slides in 3% hydrogen peroxide for 15 minutes.

3. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

4. Heat-Induced Epitope Retrieval Using The Decloaker

Add 500 ml of distilled water to the pan inside the decloaker.

Place a full rack of slides into a Tissue Tek® container with 200 ml of 1X EDTA

(Insert blank slides into any empty slots in the rack to ensure even heating of slides)

Place the container stably inside the pan and decloak for 5 minutes. *Maximum Pressure* _____

Depressurize for 10 minutes.

Remove pan top and cool for 10 minutes. *Temperature Before Cooling Slides* _____

Rinse the slides in 2 changes of distilled water for 3 minutes each time.

5. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

6. Block with the Rodent Block M Reagent for 30 minutes at room temperature.

Lot # _____ Exp. Date _____

DO NOT RINSE SLIDES WITH BUFFER BEFORE ADDING PRIMARY ANTIBODY.
ONLY WIPE EXCESS BLOCK.

7. Apply primary antibody at a 1:1000 dilution. Incubate 1 hour at room temperature.

Lot # _____ Exp. Date _____

For negative control slides, dilute mouse IgG2b control serum so that it's IgG2b protein concentration matches that of the primary antibody (if necessary). Then make a 1:1000 dilution. If the concentrations can't be matched using this method, the dilution for the negative reagent may need to be adjusted. Apply the negative and incubate for 1 hour at room temperature.

Lot # _____ Exp. Date _____

8. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

9. Apply the MM HRP-Polymer Reagent, and incubate for 20 minutes at room temperature.

Lot # _____ Date Reconstituted _____

10. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.
11. Apply the DAB chromogen. Incubate in the dark for 6 minutes at room temperature.
(Add 1 drop of DAB per ml of substrate)
Lot #_____ Exp. Date_____ New Kit: yes / no

12. Rinse the slides in tap water 3 minutes.

13. Counterstain with hematoxylin for 20 seconds.

14. Rinse the slides in tap water until water is clear.

15. Gently agitate slides in 1X wash buffer until the tissues turn blue.

16. Dehydrate through the following solutions:

Solutions	Repetitions	Time
95% Ethanol	1 time	3 minutes
100% Ethanol	3 times	3 minutes
Xylene	2 times	5 minutes

17. Coverslip

Updated 04/11/12