



## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



## Clinical Case Studies for Students and Health Professionals

The following examples are included to help students and clinicians explore in more detail the health impacts of climate change and provide real-world examples and case studies.

**Adopted and modified from:** Luber, G. and J. Lemery (2015). *Global climate change and human health: From science to practice*. John Wiley & Sons.

### Extreme Heat:

#### **Health care spending, heat and care for the elderly:**

The world's older population continues to grow at an unprecedented rate. In 2015, 8.5 percent of people worldwide (617 million) were aged 65 and older, and this percentage is projected to jump to nearly 17 percent of the world's population by 2050.<sup>i</sup> Elderly populations have unique health needs that are likely to be exacerbated with rising ambient temperature and therefore must be considered as we make projections of future health care costs.

Human mortality has been shown to increase on hot days, often attributed to cardiovascular and respiratory collapse, and the elderly are disproportionately affected for medical as well as social reasons.<sup>ii</sup> Aging is associated with a decreased physiologic ability to compensate to heat, especially when concurrent chronic and degenerative disease exists. Cognitive disability compounds these factors by altering risk perception and protective behaviors.<sup>iii</sup> The trend in increasing mortality among the elderly exists worldwide.<sup>iv</sup> With projected increases in extreme heat events and other natural disasters, the needs of the elderly are likely to be more than are projected by current estimates that do not account for climate change. Hospital administrations and clinicians must therefore prepare for this increased demand in services and curtail their systems-based practices to be able to meet the rising need with a limited financial budget.

*Climate change poses unique risks to a rapidly growing demographic of elderly patients. Clinicians, hospitals, policy makers and financial planners must prepare for these current and future needs.*

#### **Individual risk and emergency readiness:**

Epidemiological research shows that mortality in urban populations increases as the temperature rises.<sup>v</sup> Models have used mean temperature, maximum temperature, and humidity, as well as the time/rate of onset of these variables to predict when clinically significant heat waves may occur. These models allow for initiation of time-sensitive warnings to be released to the public. However, there is significant variability in the ways individuals are impacted; factors such as age, housing architecture, socioeconomics, prevalence of chronic disease, and relative social isolation all play a part. Thus, each demographic has a different threshold at which heat-related illness becomes clinically apparent. Emergency medical systems (EMS)<sup>vi</sup>, nurse lines, and emergency departments<sup>vii</sup> are at the forefront of detecting and treating heat-related illnesses. Research has shown that reliance on these organizations increases with rising temperatures, and the public's reliance on these institutions could therefore be an accurate indicator of the appearance of clinically relevant heat related disease. More research is needed to determine real-time surveillance data generated from these clinical settings could assist public health officials in deciding when to issue heat warnings to a given community. Early

## Educational Materials

A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health




warnings can help to ease the toll of heat-related illness and prevention may ease the burden of such events on the health care system.

*Integrating weather modeling and public health intervention to address vulnerable populations may ease the burden of heat stress on individuals and the health care system.*

### Heat early warning systems (HEWs): Case study: Ahmedabad, India:

Heat exposure is becoming more wide-spread and is a leading cause of weather-related morbidity and mortality. Instances of higher peak daily temperatures and longer, more intense heat waves are expected to become increasingly frequent on a global scale due to climate change.<sup>viii</sup> Adaptive measures, including the use of heat early warning systems (HEWs) and heat actions plans can safeguard the health of vulnerably populations and allow for health-care system readiness.<sup>ix</sup> In Admedabad, India following a devastating heat wave in 2010 that lead to over two thousand deaths, the Admedabad Municipal Corporation in collaboration with the Indian Institute of Public Health created a heat action plan and HEWs.<sup>x</sup> Metrologic data is used to forecast extreme heat events up to 5 days out. Established thresholds of predicted or actual temperature trigger a sequence of government and community level action. These actions include: Communication to the public through email, radio, SMS, local media and mobile applications; activation of health care system response, including increased staffing and provision of medical supplies at hospitals and health centers in affected localities; activation of designated “cooling centers” and provision of drinking water at public locations; and ongoing surveillance and assessment of public health impacts. This system is the first of its kind in India.



Left: Temperature and all-cause mortality correlation during the 2010 heat wave in Ahmedabad as compared to 2009 and 2011. Credit: [Admedabad heat action plan](#), 2016



Left: Public Health education is fundamental to heat action plans. This taxi displays a heat warning/education message distributed by the heat action committee. Photo credit: [Admedabad heat action plan](#), 2016

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### **Energy security and integrative approaches to heat stress in the developing world:**

Air-conditioning has become the mainstay approach to buffer the deleterious health effects of extreme heat events. Unfortunately, air conditioning in and of itself places major strains on energy supplies and contributes substantially to CO<sub>2</sub> emissions which in turn increase global surface temperatures. Eighty-percent of energy for air-conditioning comes from fossil fuels and according to estimates, total world air conditioning consumes roughly 1 trillion kilowatt hours annually, more than twice the total energy consumption of the entire continent of Africa.<sup>xi</sup> Given the necessity of cooling, both now and in the future, essential energy infrastructure planning must begin now. The rapidly transforming developing world contains 38 of the largest 50 cities on the planet, the warmest of which are in the developing world.<sup>xii</sup> We can anticipate that under warming conditions, the energy needed to keep these populations safe will be exponential. Thus, to curb the health effects of heat stress among vulnerable populations in a sustainable and energy-wise way, city planners and engineers must creatively utilize energy-saving technologies as well as integrate traditional technologies that have a small energy footprint. These designs include passive cooling systems (evaporative cooling<sup>xiii</sup>, night flushing<sup>xiv</sup> and passive downdraft evaporative cooling<sup>xv</sup>), exterior heat sinks, and modification of existing structures with awnings, reflective paint, and landscaping that maximizes shade. Health care organizations should model appropriate building codes and use of indoor climate control to model climate-literacy, to save costs and to achieve energy security.

*Rapid warming as well as rapid development are occurring in the developing world. To keep populations safe-guarded against the negative effects of extreme heat, innovative cooling solutions are necessary.*

### **Heat waves and primary care:**

During the 2003 heat wave in Europe, an estimated 30,000 individuals succumbed to heat death. A majority of the 30,000 deaths were among the elderly, who remained alone in their homes despite warnings to seek cooler environments.<sup>xvi</sup> The elderly are particularly vulnerable to heat waves as their mobility, hearing, vision and/or cognition may be compromised, making it difficult or impossible for them to process and/or adhere to warnings. Cardiovascular, renal and pulmonary diseases, which disproportionately affect the elderly, compound cognitive and mobility issues and thus increase vulnerability during heat waves. Primary care physicians can begin to incorporate “heat vulnerability” in their preventative health screening as a way to raise awareness among this population of the early warning signs of heat stress.

*Elderly populations are particularly vulnerable to heat stress, a factor that should be incorporated into routine care.*

### **Outdoor Air Quality:**

#### **Disparities in pediatric asthma:**

The Centers for Disease Control and Prevention report that asthma prevalence increased from 7.3% in 2001 to 8.4% in 2010, thus affecting 25.7 million individuals annually.<sup>xvii</sup> Children constitute a disproportionate majority of this statistic. It has been shown that children living in poverty experience higher rates of asthma across all ethnic groups compared to higher income households. Children of low-income families and racial minorities also manifest more severe disease than white, higher income children.<sup>xix</sup> One explanation for this finding is that children living in low-income environments are subject to substandard housing with more triggers such as mold, rodents and cockroaches. In addition, families often are forced to go without preventative medicines and heating/cooling in order to save money, which leaves children vulnerable to attacks. Deteriorating air

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



quality due to climate change is a concerning threat to children in low-income, inner-city environments. Clinicians, regulators, public health officials and housing authorities must come together to address this large and vulnerable population.

*Deteriorating air quality threatens the health of vulnerable pediatric patients, especially among those living in low-income environments.*

### **Wildfires and human health:**

Although direct impacts of wildfires tend to occur in restricted geographic regions, they have potential to impact the health of populations hundreds of miles away.<sup>xx</sup> Smoke from the combustion of biomass contains a myriad of chemicals known to be detrimental to human health.<sup>xxi</sup> Examples include carbon monoxide, organic acids, mercury and fine particulate matter. Some experts compare the impact of particulate matter produced by wild fires to that of industrial urban environments where fine particulate matter is associated with a wide-range of adverse health effects including neonatal and cardio-respiratory mortality, exacerbations of respiratory and cardiovascular conditions and pathophysiologic changes in inflammatory and coagulation pathways.<sup>xxii</sup> Carbon monoxide poisoning can lead to permanent neurologic damage. Mercury can cause neurologic toxicity and a recent study from the National Center for Atmospheric Research found that 48 tons of mercury are redistributed by wild fires annually in the United States. It is unclear how the metal is affecting human health although experts suspect it is entering waterways and drinking water.

*Smoke from wildfires can affect communities located hundreds of miles downwind with demonstrable impacts on health in the acute and likely long-term period.*

### **Ozone: a threat-multiplier:**

Acute and chronic ozone exposure has been associated with significant adverse health effects in humans, including cardiopulmonary and respiratory morbidity and premature mortality.<sup>xxiv</sup> Abrupt daily increases in ozone concentrations have been shown to decrease pulmonary function<sup>xxv</sup>, increase asthma exacerbation rates and increase emergency department visits.<sup>xxvi</sup> Ozone has also been associated with an increased relative risk of death from all cardiopulmonary causes. A recent assessment found that as many as 2,500 ozone-related premature deaths, 3 million cases of acute respiratory symptoms and one million school days could have been avoided annually could the nation attain the 75 ppb standard goal.

*Ozone not only compounds global warming but also causes measurable negative health effects during periods of acute increase.*

### **Compounding health effects of carbon-based energy:**

Fine particulate matter, generated through the burning of fossil-fuels in industrial processes and the transportation sector have been shown to cause negative acute and chronic health problems. Studies show that elevations in ambient particulate matter are associated with increases in ST-elevation myocardial infarction and accelerated atherosclerosis.<sup>xxviii</sup> Other studies show that each 10- $\mu\text{g}/\text{m}^3$  elevation in fine particulate air pollution is associated with approximately a 4%, 6%, and 8% increased risk of all-cause, cardiopulmonary, and lung cancer mortality, respectively.<sup>xxix</sup> These increases in severe, acute disease have large impacts on emergency departments and health care systems.

*Ambient fine particulate matter, produced as a byproduct of fossil fuel combustion, significantly negatively impacts human health which in turn impacts health-care system usage.*

## Educational Materials

A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health




### Extreme Weather Events:

#### Extreme weather events and chronic disease:

Extreme weather events have the potential to displace individuals from their communities and homes. Such forced movement creates disruptions in continuity of care, access to health care services and severs relationships between doctors and patients. For example, patients with conditions such as end stage renal disease often require tri-weekly dialysis in dedicated centers. In the wake of natural disasters, care for these patients and others is frequently interrupted. Individuals struggle to find new providers and centers of care while at the same time, doctors and emergency departments become overwhelmed by an influx of new patients many of whom arrive without medical records. Such interruption of treatment may contribute to the high rates of acute exacerbations of chronic diseases such as end-stage renal disease, asthma, cardiovascular disease and chronic obstructive pulmonary disease seen in the wake of natural disasters.<sup>xxxxxi</sup>

Depending on the location of a disaster and the refugee mitigation efforts, the responsibility for care of displaced patients can be spread in unpredictable ways and in far-reaching locations as was seen in the aftermath of Hurricane Katrina in 2005.



*Extreme weather events and resulting disturbances in continuity of patients with chronic medical conditions may lead to disease exacerbations. The burden of treatment is spread unpredictably among health care facilities in surrounding areas.*

Post-Hurricane Katrina distribution of refugees from 1.3 million Gulf coast households. Credit: [USA Today](#) and Federal Emergency Management Agency

#### Case study: Hurricane Sandy:

The closing of Bellevue hospital in New York City in the wake of the flooding caused by Hurricane Sandy highlights the need to intensify resources in inner-city, climate-vulnerable areas. Bellevue Hospital is the largest public hospital in the city, with 828 inpatient beds and multiple clinics that handle nearly 500,000 outpatient and 145,000 emergency visits per year. More than 80% of Bellevue patients come from the city's medically underserved populations.<sup>xxxii</sup> When Superstorm Sandy hit New York on the evening of October 29, 2012, it caused power-outage throughout the hospital and severe flooding which ultimately forced the hospital to evacuate all patients and close its doors. To make matters worse for patients, the hospital remained closed for three months after the storm. The effect that this closure had on city's most vulnerable populations is unknown. Those who relied upon Bellevue for intensive outpatient care or those requiring hospitalization would have had to travel outside their neighborhood on unreliable post-disaster public transportation to seek care from other already severely over-crowded hospitals. In retrospect, there were many aspects of Bellevue's design that were not prepared for a storm surge, despite its location along the water. This incident highlights the need to intensify resources in areas such as inner cities, which have high concentrations of at-risk populations and are vulnerable to natural disasters.

*System-wide evaluation of public and "safety net" hospitals is needed to prepare for natural disasters.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### Health-related exposures from heavy precipitation events:

Expanding areas of urbanization across the United States coupled with increasing extreme precipitation events pose significant threats to human health. During times of low precipitation, urban environments collect substances such as heavy metals, chemical pollutants, and pathogens on their impervious surfaces. When extreme precipitation occurs, treatment capacities of water facilities are overwhelmed, and these chemicals and pathogens are released into surface, drinking, and recreational waters. It is estimated that 6%-40% of the 99 million cases of acute gastroenteritis that occur annually in the US can be attributed to contaminated drinking water. We know that more than half of contaminated drinking water outbreaks in the past sixty years have followed extreme rainfall.<sup>xxxv</sup> In addition to bacterial exposures, toxic substances such as copper, zinc, and lead, as well as pesticides and hormonally active compounds accumulate on roads, roofs, and parking lots and are released into the drinking and surface waters during extreme precipitation events, never even seeing a treatment facility. These substances have the propensity to create neurotoxicity and act as carcinogens when ingested. The individuals most at risk from these types of environmental contamination include children, the elderly, pregnant women and the immunocompromised. Physical factors that put communities at risk can be measured through indices such as the normalized built-up difference index (NDBI) and thus be used to target improvements in domestic water management.

*Extreme precipitation events unleash toxic man-made compounds as well as infectious pathogens into drinking and recreational waters, posing significant risks to human health.*

### Flooding and environmental contamination:

Severe weather events are associated with flooding and release of toxins from contained sources that have the potential to enter the human food chain and affect food safety. For example, following the 2002 floods in central Europe, monitoring programs traced polychlorinated dibenzo-p-dioxins and dibenzofuranes from breached containers into soils and into the food chain via cow milk.<sup>xxxvii</sup> Floodwaters from Hurricane Katrina caused oil spillage from storage tanks as well as spillage of pesticides, metals and stored hazardous waste that were untraced and thus their impact unknown. It is unclear what the significance these toxins is to human health, however it is a field that deserves further monitoring and intervention in order to ensure a safe food supply especially during disasters.

*Contained industrial wastes are susceptible to being unleashed during extreme weather and flooding leading to release of toxins that have the potential to enter the food chain and adversely affect human health.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### Health Care Facility Challenges and Opportunities:

#### **Healthcare Facilities**

During natural disasters, hospitals are considered “custodial” organizations and thus are the last to be evacuated and closed. Health facilities can be significantly impacted during tropical storms. Flood impacts and landslides due to extreme precipitation can damage backup generators in basements, leading to no electricity, running water, sewage treatment, refrigeration or access to CT scanners, X-rays, laboratory information, or blood banks. Ventilators and dialysis machines would not function and essential supplies can dwindle due to transportation breakdown. Many hospitals that are short-staffed have providers working long shifts in adverse conditions witnessing the inevitable deaths of patients who would not be sustained without life support.<sup>xxxix</sup> How should we prepare for sudden or prolonged resource shortage? Over the past fifty years, modern medical practice has become increasingly dependent upon technology in diagnostics and treatment, and health care is one of the most “energy intense” sectors of the economy.<sup>xl</sup> Hospitals can prepare for resource shortage by adopting green, energy saving practices, by identifying through institutional analysis, what essential systems are necessary in emergency scenarios and how to shunt limited power to those areas. Health care providers can and must begin to train themselves to practice medicine without intensive technology or essential diagnostic testing. This approach will not only prepare us for disasters related to the environment but also abate the current financial healthcare crisis.

*Health care is currently one of the most energy-intensive sectors and its systems become vulnerable during natural disasters when resources become scarce. Hospitals and providers can prepare by adopting green infrastructure and limiting unnecessary testing and interventions in the present.*

#### **Case Study: Tropical Storm Erika**

In August 2015, Tropical Storm Erika struck Dominica, causing at least 31 deaths. 80% of the island was left without power and 14,291 people were rendered homeless.<sup>xii</sup> The combination of intense rainfall, unusual dry season and cracking of clay soils contributed to slope failures and debris generation resulting in major damages in Dominica.<sup>xiii</sup> Several health care facilities were significantly affected either directly by flooding or because of isolation from the rest of the country through road blocks from landslides, roadways being damaged, or bridges being washed away. Nine primary healthcare facilities were closed for at least two days and many were blocked from vehicle access requiring airlifts for patients. 10 of 53 health facilities did not have water connection and 14 did not have solid waste collection after the storm. Healthcare facility damage costs were estimated at more than \$1,727,900. Water supply became an urgent need in many of the health facilities and a focus of disaster response efforts.

*Read addendum for more information.*

### Vector-Borne Diseases:

#### **Preparing for change: Malaria early warning systems**

The transmission of malaria through *Anopheles* mosquitoes is strongly linked to climate conditions. Predicting epidemic outbreaks of malaria in relation to environmental conditions has therefore become an area of intense research. Malaria early warning systems combine elements of early detection of epidemics (case surveillance), early warning (based on monitoring meteorological data) and long-range climate modeling.<sup>xiv</sup> As opposed to endemic malaria, which occurs in populations that have some immune resistance to the parasite, epidemics occurring in immune-naïve communities have more disastrous health consequences with higher rates of morbidity and mortality. The health

## Educational Materials

A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health



impacts of accurate prediction of epidemics appear promising. A recent model developed by Thompson and colleagues is able to predict epidemic occurrence with reasonable certainty with greater than one month and at times five-month lead time.<sup>xiv</sup> Advocates for early warning systems urge that precise prediction models give policy makers and health officials the time necessary to protect vulnerable populations in proactive versus reactive way. The question now is whether the models are sound enough to put into practice and if policy makers are willing to respond.

*Malaria early warning systems combine case surveillance with long-range climate modeling to allow clinicians and policy makers to proactively prepare for outbreaks and allocate resources.*


### Forecasting climate-sensitive emerging diseases – Rift Valley River:

Rift valley fever (RVF) is a mosquito transmitted viral disease that affects ruminant animals and may spread into humans through direct contact with the blood or organs of infected animals.<sup>xlvi</sup> The virus was first identified in 1931 in Kenya, and since then more than 30 outbreaks have occurred in Africa. Until 2000, the disease was confined to the African continent. However, in 2000, an outbreak was identified in Saudi Arabia with 683 laboratory-confirmed human cases and unknown numbers of affected livestock. Outbreaks are highly lethal in animal populations, leading significant economic losses and hunger in rural and nomadic populations. The 2006-2007 RVF outbreak in East Africa is estimated to have cost \$32 million to Kenya alone from livestock losses and export bans.<sup>xlix</sup> In humans, clinical manifestations range from a mild acute febrile illness to severe encephalopathy, liver failure, disseminated intravascular coagulation and death.<sup>l</sup>

Outbreaks of RVF are strongly connected to climatic conditions, most importantly the El Nino/Southern Oscillation, and the resulting patterns of heavy rainfall that occur over normally arid geographic regions of East Africa approximately every 10 years.<sup>li</sup> Heavy rains create areas of standing water in grassland depressions, termed “dambos,” within land used to graze domestic animals. These pools serve as ideal breeding habitat for the Culex and Aedes mosquito and the nidus for outbreaks of RVF.<sup>liiiiii</sup>

The close connection between sub-seasonal climate variations and disease outbreaks have enabled the successful development of forecasting models and early warning systems that provide essential lead-time for public health authorities to take measures to reduce negative societal, economic and health impacts.<sup>liv</sup> Currently, a collaboration between NASA, CDC, USDA and DOD issues notification of emerging threats of RVF to at-risk countries based on climate models.<sup>lv</sup> In regards to the outbreak in Saudi Arabia, experts hypothesize that the virus was introduced into the Arabian peninsula during the 1997-1998 epidemic in East Africa via infected mosquitos or infected livestock and that climatic conditions facilitated the outbreak.<sup>lvi</sup> The spread of RVF into non-endemic regions highlights the urgent need to enhance vector surveillance and strengthen collaboration and coordination across international, public health, agricultural and meteorological sectors to develop and test reliable climate-disease forecasting tools.

**Right:** Risk forecasting map for Rift Valley Fever East Africa. Dark Green: known or predicted presence of RVF. Dark red: areas with recent heavy rainfall. (Photo Credit: [NOAA](http://NOAA))



## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### Human migration and vector borne illness

Climate change is likely to cause mass movements of people - from rural areas into urban zones, as well as migrations across country lines in response to changes in resource availability. Therefore, clinicians world-wide must have a high index of suspicion for the appearance of non-endemic diseases. For example, the West Nile virus first made its appearance in Uganda in 1937 and then spread to Tunisia and Israel before appearing in New York City. The spread of this virus, as well as others such as Ebola and avian influenza A (H5N1) exemplify the interconnectedness of our environment and highlight the need for international communication about the appearance of infectious vector-borne diseases.

*The impacts of climate change on the geographic distribution of resources may result in mass human movement. Clinicians and public health officials must be alert to the possibility of infectious diseases appearing in unexpected places.*

### Climate and Zoonotic disease transmission

Zoonotic pathogens account for 61% of all infectious pathogens and 60% of emerging infectious diseases.<sup>lvii</sup> Anthropogenic land use patterns, climate change, hydrologic cycle changes and urbanization influence transmission of these pathogens by altering ecologic niches of the vector and host, by changing community structure, and by altering behavior of vectors, hosts and humans. Such complexity makes it difficult to predict whether changing climates will lead to increased or decreased disease incidence.<sup>lix</sup> What we do know is that immunology and physiology play a large role in determining the severity of infection as well as its further transmission. Poor nutrition, environmental stress and prior immunologic exposure all contribute to the robustness of the human host's defense and determine parasitic, bacterial and viremic loads and thus how contagious a patient will become.<sup>lx</sup> Malnutrition and chronic diseases are widespread in both the industrialized and developing world and thus play an integral part in the spread of zoonotic pathogens.

*The manifestation of disease from climate sensitive zoonotic pathogens is influenced by the baseline health of communities and individuals which may in turn be threatened by climate change.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### Water-Related Infection:

#### **Climate and Cholera:**

We are currently in the midst of the seventh global cholera pandemic, which began in Indonesia in 1961. However, unlike the prior six pandemics which “burned out” after 5-20 years, the current strain of *Vibrio cholerae* O1 has undergone two serotype changes<sup>lxi</sup> and persists to the present day. It remains endemic in 50 countries, and re-emerges catastrophically during natural disasters, civil unrest and the breakdown of public health measures. Examples are evident in the Rwandan<sup>lxii</sup> refugee camps in 1994 and in 2010 following the earthquakes in Haiti.

Recently, the current pandemic strain of *Vibrio cholerae*, El Tor, has undergone a hybridization with the classic serotype and become significantly more virulent, killing 1-5% of its hosts as compared to traditional rates of less than 1%.<sup>lxiv</sup> There are currently two killed oral vaccines approved for use, Dukoral® and Shanchol®. Both require multiple doses and provide a limited time frame of immunity. The World Health Organization does not currently recommend routine use of the vaccines in the immediate outbreak of an epidemic, as data is insufficient to predict whether immunity is conferred fast enough to curb the spread of disease, and the cost of administration is high. It instead recommends focusing efforts on providing clean drinking water and rehydration as primary management strategies.



*Climate change is likely to intensify preexisting public health emergencies such as cholera. Increase surveillance is needed in the setting of climate related disasters given the ongoing threat of this global pathogen.*

Left: *Vibrio cholerae* (Photo credit: [Wikipedia](#))

#### **Impacts of cryptosporidium on immunocompromised patients:**

Cryptosporidium is an ubiquitous and tenacious organism that is highly resistant to conventional means of disinfection (chlorination and filtration) owing to its small size (4-6 micrometers) and cystic structure.<sup>lxv</sup> Despite even stringent standards for water turbidity in wastewater treatment facilities, outbreaks still frequently occur.<sup>lxvi</sup>

In developing countries *Cryptosporidium* accounts for 6.1% of cases of diarrhea in HIV-negative patients and 24% of cases in HIV-positive patients, while in developed countries it causes 2.1% and 13.8% of illnesses respectively.<sup>lxvii</sup> In the immunocompromised population it can often be a fatal disease. Even when not fatal, infection may also cause dire yet subtle complications. Among children living in impoverished areas, infection has been shown to lead to impaired weight gain in the month following symptomatic and even asymptomatic infection, which occurs without a subsequent “catch up” growth period.<sup>lxix</sup> Among patients with chronic diseases, infection has been observed to interfere with the absorption and therapeutic levels of anti-retroviral and anti-tuberculous drugs,<sup>lxx</sup> which can affect the transmission of these highly communicable diseases.

*Cryptosporidium is a common climate-sensitive disease that health care providers must be aware of in order to prevent complications among vulnerable patients.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### How safe is our water supply?

In 2012, the World Health Organization announced attainment of the Millennium Development Goal (MDG) of reducing the number of individuals without “sustainable access to safe drinking water” by half. One major information gap that remains regards the *quality* of the water. Even in the best of public water treatment scenarios, outbreaks of water-related disease still occur.<sup>lxxi</sup> In a recent study conducted by the UNICEF and USAID, households were questioned in regards to their drinking water access. As a proxy for “sustainable access to safe drinking water,” individuals were asked if they had “use of an improved water source,” where “improved” was generalized to imply only a mechanism of handling water that aimed to decrease the likelihood of it causing disease or infection. Since there is no agreed definition of “safe” and no widely used and inexpensive means of testing drinking water for pathogens, it is difficult to predict if attainment of this MDG will result in a decrease in the global burden of gastrointestinal disease. In addition, we are at this point unable to predict with regional certainty the effects of climate change on local water access. Medical personnel will likely be at the forefront of monitoring this impact.

*Access to safe water is a difficult milestone to achieve, define and assess. It is requisite to achieve modern health standards and may be threatened by changing climates.*

### Health implications of Harmful Algae Blooms (HAB):

Climate change is leading to more frequent and widespread HABs thus increasing the incidence of human exposures. The impacts of HABs are diverse and may affect both coastal and inland communities. For example, *Kareina brevis*, the dinoflagellate responsible for Florida’s red tide,<sup>lxxiii</sup> blooms on an annual basis with varying degrees of intensity. The cells of these organisms contain potent neurotoxins, which when lysed by wind and waves become aerosolized and can travel up to a mile inland.<sup>lxxv</sup> Studies have linked the blooming of these algae with upper and lower respiratory symptoms including rhinorrhea, cough, and bronchoconstriction, especially in asthmatics and the elderly. High intensity blooming events have been shown to correlate with increased emergency department usage in nearby coastal areas. Climate change and resulting changes to oceans may increase the intensity and occurrence of HAB’s, the costs of which include exacerbation of chronic illness, and increased volumes of patients in emergency departments.

*Harmful algae blooms pose significant risks to communities and health care systems. Strong public health monitoring is needed to trace these impacts and create notification pathways within the health care system and within communities.*

### Food and Nutrition related impacts:

#### Micro-finance, nutrition and women’s empowerment in the era of climate change:

Climate change is likely to alter the price and availability of food and thus impact nutrition and health.<sup>lxxvii</sup> Resource-poor communities and those whose livelihoods depend on agriculture are most susceptible to food scarcity. According to the German Development Institute, women produce between 60 and 80 percent of all food in the developing world and thus represent a highly vulnerable group. Additionally, women are prone to nutritional deficiencies secondary to increased needs during pregnancy and nursing and these matters can be intensified by household food hierarchies. Undernutrition has profound effects on neonatal development and is associated with intrauterine growth restriction, pregnancy complications and perinatal mortality. When resources are scarce within a household or community, the decision of how and when to allocate finances is crucial.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



Global reduction of gender inequality and food insecurity are paramount to the United Nations Sustainable Development Goals. Gender inequalities include impaired access to education and healthcare as well as limited power to influence household financial decision-making. Indexes have been developed to quantify gender disparities in household power.<sup>lxxxix</sup> Evidence suggests that when women have higher levels of household power, the overall health of children improves, likely due changes in allocation of financial resources towards purchasing food and healthcare. Empowering women to establish businesses and manage finances through micro-loan programs in have shown that participation in loan programs is directly correlated to increases in women's household power.<sup>lxxx</sup> Thus micro-finance directed towards women may positively affect household resource allocation and health, thereby building resilience in communities most vulnerable climate shocks.

*Financially empowering women builds resilience in communities at risk of climate shocks and resulting resource and nutrition scarcity.*

### **Amplification of repercussions from infection on nutrition:**

Experts estimates that one third of the world population is infected with some form of intestinal helminth parasite at all times.<sup>lxxxii</sup> Helminth infections have been correlated with impaired physical and cognitive growth and development in children. *Ascuris lumbricoides*, *Trichuris trichiuru*, *Schistosoma* species, and other hookworms colonize the intestinal lining. Colonization prevents the absorption of nutrients and causes a chronic inflammatory state, diarrhea, and blood-loss anemia. Helminth life cycles involve periods of dormancy in the environment and are likely to be impacted by changing global climates. Researchers propose that either the infection itself or the secondary manifestations of malnutrition, stunting, and anemia have effects on cognition and learning among infected children

Recently, studies looking at the effects of improved nutrition suggest that dietary intake of micronutrients such as vitamin A, vitamin B12, vitamin C, riboflavin, zinc, selenium, and iron all have immunomodulating functions and may impact the course of parasitic disease. Thus, a positive feedback situation emerges whereby dietary malnutrition increases risk for parasitic infection, which itself furthers the extent of the malnutrition through malabsorption. A new clinical approach towards eradication of infection could focus on intervening in micronutrient deficiencies while at the same time employing the use of chemotherapeutic agents and public health measures.

*Helminth infections are a prevalent climate sensitive illness that has wide-reaching implications for human health and nutrition.*

### **Food scarcity and agrochemicals:**

Climate change and population growth are increasing demands on the global food supply and increasing reliance upon industrialized food production and agricultural chemical use. Research into the long-term health effects of pesticide use lags decades behind widespread utilization of these compounds. Large studies of the chronic effects of pesticide exposure on children's health have shown links to birth defects, premature birth, intrauterine growth restriction and several childhood cancers.<sup>lxxxviii</sup> Many pesticides currently in use are classified as carcinogens by the United States Environmental Protection Agency. Expert consensus now holds that there are links between pesticide exposure and neuro-cognitive development. Prospective studies link early-life exposure to reductions in intelligence quotient and increased rates of attention-deficit/hyperactivity disorder and autism.<sup>xcxci</sup> Unfortunately, there is currently no information on developmental toxicity of over 80% of

## Educational Materials

*A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health*



the 3000 chemicals used in US agriculture.<sup>xcii</sup> And yet neurodevelopment disorders such as autism, attention-deficit disorder and pervasive development disorder now affect one in six children living in the industrial world. More research is necessary to untangle the myriad of connections between early life exposure and pesticide toxicity and create regulations that address the severity of the clinical implications of exposure to ensure a safe food supply as we rely more upon industrialized food.

*Climate change leads to food insecurity, more reliance on industrialized food production and the use of chemicals with unknown and known detrimental effects on human health.*

### **Climate preparedness: Food Security**

Climate change is anticipated to have far reaching impacts on food production and certain aspects of food quality.<sup>xciv</sup> Currently, one-third of children in developing countries are malnourished by anthropometric standards. It is essential use an evidence-based approach to understand why current systems are failing as researchers and policy makers attempt to stabilize food security. Central America has the highest prevalence of stunting (23%) and the lowest rates of improvement over the past 25 years, while nearby South America has lower levels stunting and the highest rates of improvement, despite economic stagnation in both regions.<sup>xcv</sup> A recent Lancet series analyzes how and why nutrition programs fail, and reports that some failures can be traced to a lack of political commitment, while others can be traced to failure to support programs that have evidence-based results.<sup>xcvi</sup> What public health professionals and clinicians can learn from this is that malnutrition is knowable and treatable and that with proper programs, we have the tools to address the issue. Clinicians must serve as spokespersons in their local political arena to advocate for evidence-based nutrition programs.

*Evidence-based approaches should be used to assess and address the health impacts of food security among vulnerable populations.*

### **Climate change and food scarcity:**

Global climate change is forecasted to intensify global food shortages in vulnerable places. The effects of chronic undernutrition range from physical to cognitive. On a worldwide level, growth stunting, severe wasting and intrauterine growth retardation together contribute to 2.2 million deaths annually, accounting for 35% of mortality of children less than five years of age.<sup>xcvii</sup> Malnutrition has been shown to lead to impaired cognitive development, learning disability, deferment of education, attrition from the work force and less decreased lifetime earnings. Micronutrient deficiencies are pervasive as well and are likely to intensify. Iodine deficiencies result in goiter, hypothyroidism, and developmental disabilities including severe mental retardation. Insufficient intake of vitamin A is the leading cause of blindness worldwide. Dietary iron deficiency leading to anemia is worsened by chronic parasitic infection and chronic illness that so often accompanies it in resource-poor settings. Folic acid is essential for normal nervous system development, yet is often lacking in diets along with zinc, and both are linked to childhood growth stunting.<sup>xcix</sup>

*Health effects of chronic undernutrition manifest in a myriad of ways, ranging from physical to cognitive, and are likely to intensify in the future as a result of climate change.*

### **Ciguatoxins: health effects of a climate sensitive toxin**

Ciguatera toxin is one of the most common non-bacterial causes of fish-related food poisoning in the United States.<sup>c</sup> The disease is caused by consumption of certain tropical reef fish which bioaccumulate toxins from the algae *Gambierdiscus toxicus*. Although most cases are limited to the algae's geographic range of Florida and Hawaii, changing marine climates could have unpredictable effects on algae

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



populations and cases of disease could rise. In addition, specialty fish are often shipped around the country causing the disease to manifest outside its endemic range. Therefore, it is important for health care providers to be able to recognize the symptoms of poisoning. Ciguatoxin is a heat and acid stable, odorless, tasteless<sup>ci</sup> compound that causes hyperpolarization of sodium channels in an array of tissues,<sup>cii</sup> leading to a broad array of neurologic symptoms. Common manifestations include paresthesias, numbness, headache, ataxia, vertigo, weakness, paralysis and cranial nerve dysfunction.<sup>ciii</sup> Gastrointestinal symptoms and hemodynamic instability also occur. Treatment is supportive, yet effects may last weeks. The differential diagnosis for Ciguatoxin poisoning includes paralytic shellfish poisoning, stroke, eosinophilic meningitis, organophosphate poisoning, tetrodotoxin poisoning and more. Clinicians must take a careful history to uncover this rare yet important cause of climate sensitive bio-poisoning.

*Ciguatera poisoning is a climate sensitive cause of bio-poisoning that clinicians must be aware of.*

### **The global nutrition transition, “dual-burden disease” households, and climate change:**

Global climate change directly impacts agriculture and the cost and availability of food. Stark differences exist between rich and poor nations in terms of the prevalence of various diseases; communicable diseases and malnutrition predominate in developing countries whereas non-communicable diseases such as diabetes, cancer, and cardiovascular disease are more common in developed countries.<sup>civ</sup> However, this phenomenon is rapidly changing in low to middle income countries, where there is now a trend towards the existence of both types of disease within the *same household*.<sup>cv</sup> Malnourished children live alongside mothers who suffer from obesity, diabetes and cardiovascular disease, causing a “dual-burden.” The current nutrition transition is a shift from a diet rich in legumes, vegetables and coarse grains, which have low carbon-foot print, to a diet high in refined carbohydrates, sugars, fats, and animal-source foods. The forces shaping this transition are thought to be food prices, demographic shifts from rural to urban life-styles, the commercialization and packaging of food, as well as life-style changes in developing regions.<sup>cvi</sup> Researchers forecast that the impact of this transition will be epidemic levels of obesity throughout the world. From 1988 to 1998 alone, the prevalence of obesity increased from 2.3% to 19.6%.<sup>cvii</sup> Estimate predicts that by 2030 there will be approximately 2.16 billion overweight and 1.12 billion obese adults. The obesity epidemic will take a tremendous toll on health care systems in transitioning countries. Providers will have to shift their expertise and resources towards caring for the co-morbidities associated with obesity, such as diabetes and cardiovascular disease, while at the same time addressing the needs of those suffering from diseases related to communicable diseases and malnutrition.

*Global climate change is affecting food availability and lifestyle choices, thus shifting the prevalence of diseases worldwide.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### Mental Health and Well-being:

#### **Climate and mental health and recovery:**

According to the World Health Organization (2011), mental, neurologic and substance use disorders account for 14% of the global burden of disease. Current estimates place the global shortage of mental health care workers at 1.8 million, including 55,000 psychiatrists, 628,000 nurses and 493,000 psychosocial care providers.<sup>cxix</sup> The rippling effect of mental distress caused by natural disasters can be observed in increased rates of substance abuse,<sup>cx</sup> post-traumatic stress disorder,<sup>cxii</sup> child abuse<sup>cxiii</sup> and suicide. These conditions have the potential to undermine the social fabric of a community in which a natural disaster has occurred, impairing reconstruction efforts and economic resolution. Additionally, these mental health disorders may further manifest as disease through their protean interactions with physical health.<sup>cxiv</sup> Although regions of the world with a disproportionate dearth of trained mental health work force are likely to experience a greater impact, industrialized nations are not immune. Currently, the United States spends a relatively low percentage of total health care dollars on mental health. Studies have identified a widespread prescribing and non-prescribing provider shortage,<sup>cxv</sup> which has resulted in “defacto” mental health care, whereby roughly primary care doctors or sub-specialty practitioners care for 50% of patients seeking treatment for mental health disorders. Thus, as the mental health burden from natural disasters rises, all areas of medicine are going to be called upon to care for patients in need, with little funding or systems in place to do so.

*Mental health care plays a crucial role in disaster recovery and the dearth of providers domestically and globally means the care will be distributed among all specialties of medicine.*

#### **Trauma and mental health – increasing surveillance of disaster impacts:**

Natural disasters can result in significant physical and emotional trauma. The symptoms of trauma can be expressed in many forms, which may complicate accurate clinical diagnosis and compound existing medical co-morbidities, especially chronic pain. Some experts in the field of trauma have begun to unravel the ways in which culture informs the expression of psychological distress. The Harvard Program for Refugee Trauma has employed behavioral health specialists to design unique screening tools for six different countries in order to detect post-traumatic stress disorder (PTSD) in various cultures. They have found in many cases that neurologic, somatic and visceral pains become physical manifestations of psychological trauma. The severe pain that patients report is a real experience for them, regardless of any “organic” cause recognized by western medicine. Experts point out that a large barrier to patient treatment is the fact that health care professionals often shy away from asking patients about their history of trauma because they “believe they won't have the tools or the time to help survivors once they've elicited their history.”<sup>cxvi</sup> However, this history is imperative and will possibly become more so in the future. Recent studies through the Veteran’s Administration have shown value of proper identification and treatment of PTSD. It is estimated that up to 70% of veterans that live with chronic pain have PTSD, and up to 80% of those with PTSD suffer from refractory chronic pain. Research has shown that treatment of PTSD statistically lowers a patient’s perception of pain, thus improving quality of life and decreasing dependence upon narcotic pain medications. To prepare for the future and to better handle the already heavy burden of trauma in our society, medical schools and professional organizations can design curricula to incorporate practical education on detecting, discussing and managing trauma.

*Climate change has the potential to result in widespread physical and psychological trauma. Thus, clinicians must be well versed in properly diagnosing and treating stress syndromes.*

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



### **Stress and physical health: Implications for current and future generations:**

“Stress” is a common complaint in our society. It may be conceived of as the passing of a mental, emotional or physiological threshold, after which a person can no longer cope and adapt.<sup>cxix</sup> On a physiologic level, stress is mediated through the production of the hormone cortisol in the adrenal glands. Higher levels of cortisol have been correlated with poor physical health, manifested through decreased immune function, high blood pressure<sup>cxx</sup>, reproductive disturbances<sup>cxxi</sup> and more. In an era of global climate change, stress levels are expected to be rising due to the real and perceived threats of drought, famine, severe weather, over-population, forced migration, and more. As an example, in a study of Kenyan farmers subject to drought conditions, cortisol levels were found to be significantly elevated. In another study, unemployment and job insecurity were found to impact the neuroendocrine and immune systems. Even perceived stress may have negative health implications. In one study, participants who reported that stress was impacting their health were found to be twice as likely to suffer a heart attack as those who believed that stress had no effect on their physical wellbeing. Higher stress levels have implications for current as well as future generations. For example, women living in war-torn regions of the Democratic Republic of the Congo experience high levels of stress during pregnancy that may directly influence birth weight and lead to epigenetic modification of the glucocorticoid receptor NR3C1 of the infant.<sup>cxv</sup> These epigenetic changes could impact how the child responds to stress and thus impact their long-term health profile. Clinicians will be faced with these health implications of stress at all levels of care and must learn how to advise patients.

*Climate change causes real and perceived threats of food scarcity, job insecurity, natural disasters and more. Stress has wide-reaching effects on health that clinicians must be able to promptly and accurately diagnose and treat.*

### **Recovering the psychologic fabric of a community:**

When major disasters create psychosocial trauma, individuals and relational structures among communities and families have the potential to become strained. Research has examined the prevalence of stress disorders following disasters and found that not all individuals are affected equally. While many experience acute stress immediately following events, only some go on to develop long-term and debilitating conditions.<sup>cxvii</sup> Furthermore, despite the negative connotations of stress, some investigations have found that when individuals are able to adapt after trauma, that new opportunities, deeper compassion for others, and strengthened spirituality can be the rewards. What factors contribute to an individual’s ability to adapt and evolve? Researchers have found that communication and trusting relationships may counters feelings of helplessness and meaninglessness in the setting of trauma. Therefore, perhaps disaster management must encompass mechanisms to nurture stronger community ties as a way to instill resilience. Landau and Saul outline the following helpful themes for building community in the wake of a disaster: (1) Enhance social connectedness and information and resource sharing (2) encourage involvement in collective storytelling and validation of the trauma and response with the goal of “shared meaning” (3) re-establish the rhythms and routines of life and engage in collective healing rituals (4) arrive at a positive vision of the future with renewed hope. Healers and leaders can work within their communities in order to foster these processes at many levels by raising awareness, arranging community meetings, and nurturing an open and compassionate attitude towards distress in others.

*Mitigation of climate-related disasters must consider the psychological effects that trauma has on communities. Clinicians and other community leaders can serve as leaders in rehabilitation efforts.*

## Educational Materials

*A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health*



### Addendum

#### **Case Study: Tropical Storm Erika**

Interruptions in healthcare delivery in the aftermath of climate-related disasters can create dire conditions for individuals seeking care for both acute and chronic medical conditions. Investment in resilient healthcare infrastructure, including consideration of structural and non-structural factors of health care systems, is crucial to prevent unnecessary morbidity. Structural factors include: hospital design, backup provision of clean water and electricity, energy generation, disposal of hazardous wastes and more. Non-structural and operational factors include: provision of essential supplies during disaster scenarios (medications, equipment), transportation/evacuation plans for patients and staff, coordination of patient care between regional and local health care facilities and protection, awareness and coordination of care for vulnerable populations in the community (e.g. nursing homes, emergency shelters). Initiatives and upgrades to increase the resilience of health care facilities can be paired with energy efficiency measures which have the co-benefits of reducing operating costs in addition to mitigating greenhouse gas emissions.<sup>CXXX</sup>

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



## References

- <sup>i</sup>Wan He, Daniel Goodkind, and Paul Kowal U.S. Census Bureau, International Population Reports, P95/16-1, An Aging World: 2015, U.S. Government Publishing Office, Washington, DC, 2016.
- <sup>ii</sup> Davis R, Knappenberger P, Michaels P, Novicoff W. (2003) Changing Heat-Related Mortality in the United States. *Environmental Health Perspectives*, 111(14), 1712-1718.
- <sup>iii</sup> Astrom D, Forsberg, B, Rocklov J. (2011) Heat wave impact on morbidity and mortality in the elderly population: A review of recent studies. *Maturitas*, 69, 99-105.
- <sup>iv</sup> Astrom D, Forsberg, B, Rocklov J. (2011) Heat wave impact on morbidity and mortality in the elderly population: A review of recent studies. *Maturitas*, 69, 99-105.
- <sup>v</sup> Bassil K, Cole D, Moineddin R, Lou W, Craig A, Schwartz B, Rea E. (2011) The relationship between temperature and ambulance response calls for heat-related illness in Toronto, Ontario, 2005. *J Epidemiol Community Health*, 65, 829-831.
- <sup>vi</sup> Leonardi GS, Hajat S, Kovats RS, et al. (2006) Syndromic surveillance use to detect the early effects of heat-waves: an analysis of NHS direct data in England. *Soz Praventiv Med*, 51, 194–201.
- <sup>vii</sup> Claessens YE, Taupin P, Kierzek G, et al. (2006) How emergency departments might alert for pre-hospital heat-related excess mortality? *Crit Care*, 10, R156.
- <sup>viii</sup> IPCC, “Special Report on Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation,” March 8, 2012: <http://www.ipcc-wg2.gov/SREX/>.
- <sup>ix</sup> Hess JJ, Ebi KL. Iterative management of heat early warning systems in a changing climate. *Annals of the New York Academy of Sciences*. 2016 Oct 1;1382(1):21-30.
- <sup>x</sup> AHAP (2015) Ahmedabad Heat Action Plan. Amdavad Municipal Corporation, Government of Gujarat. <http://www.nrdc.org/international/india/extreme-heat-preparedness/files/ahmedabad-heat-action-plan.pdf>. Accessed 3/12/17
- <sup>xi</sup> Dahl R. (2013) Cooling Concepts: Alternatives to air-conditioning for a warm world. *Environmental Health Perspectives*, 1(1).
- <sup>xii</sup> Sivak M. (2009) Potential energy demand for cooling in the 50 largest metropolitan areas of the world: implications for developing countries. *Energy Policy*, 37(4),1382-1384.
- <sup>xiii</sup> Evaporative Cooling: How an Evaporative Cooler Works [website]. Sacramento, CA:Consumer Energy Center, California Energy Commission (2012). Available: [http://www.consumerenergycenter.org/home/heating\\_cooling/evaporative.html](http://www.consumerenergycenter.org/home/heating_cooling/evaporative.html) [accessed 11 Dec 2012].
- <sup>xiv</sup> Night-Purge Ventilation [website]. San Rafael, CA:Autodesk, Inc. (2012). Available: <http://sustainabilityworkshop.autodesk.com/fundamentals/night-purge-ventilation> [accessed 11 Dec 2012].
- <sup>xv</sup> Kamal MA. (2011) Reinventing traditional systems for sustainable built environment: an overview of passive downdraught evaporative cooling technique for energy conservation. *J Res Arch Plan*, 11(2), 56-62.
- <sup>xvi</sup> Ledrans M, Pirard P, Tillaunt H, et al. (2004) The heat wave of August 2003: what happened? *Rev Prat*, 54(12), 1289-97.
- <sup>xvii</sup> Akinbami LJ, Moorman JE, Bailey C, et al. (2012) Trends in asthma prevalence, health care use and mortality in the United States, 2001-2010. *NCHS Data Brief*. No 94.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



- <sup>xxiii</sup> Weitzman M, Gortmaker S, Sobol A. (1990) Racial, social, and environmental risks for childhood asthma. *Am J Dis Chil*, 144, 1189–94.
- <sup>xxix</sup> Murphy JS and Sandel MT. (2011) Asthma and social justice: how to get remediation done. *Am J Prev Med*, 41(2S1), S57-S58.
- <sup>xx</sup> Kinney PL (2008) Climate change, air quality, and human health. *Am J Prev Med*, 35(3), 450-467.
- <sup>xxi</sup> N Naeher LP, Brauer M, Lipsett M, Zelikoff JT, Simpson CD, Koenig JQ, et al. (2007) Woodsmoke health effects: a review. *Inhal Toxicol*, 19(1), 67–106.
- <sup>xxii</sup> P Pope CA, III, Dockery DW. (2006) Health effects of fine particulate air pollution: lines that connect. *J Air Waste Manag Assoc*, 56(6), 709–742.
- <sup>xxiii</sup> Lipsher S. “Wildfire smoke a culprit in mercury’s spread.” *The Denver Post*. October 19, 2007. Print.
- <sup>xxiv</sup>,<sup>5</sup> Berman JD, Fann N, Hollingsworth JW, Pinkerton KE, Rom WN, Szema AM, et al. (2012) Health benefits from large-scale ozone reduction in the United States. *Environ Health Perspect*, 120, 1404–1410.
- <sup>xxv</sup> Mudway IS, Kelly FJ. (2000) Ozone and the lung: a sensitive issue. *Mol Aspects Med*, 21, 1–48.
- <sup>xxvi</sup> Choi M, Curriero FC, Johantgen M, Mills MEC, Sattler B, Lipscomb J. (2011) Association between ozone and emergency department visits: an ecological study. *Int J Environ Health Res*, 21, 201–221.
- <sup>xxvii</sup> Jerrett M, Burnett RT, Pope CA, III, Ito K, Thurston G, Krewski D, et al. (2009) Long-term ozone exposure and mortality. *N Engl J Med*, 360, 1085–1095.
- <sup>xxviii</sup> Zhang Z, Whitsel E, Quibrera PM, Smith R, et al. (2009) Ambient Fine Particulate Matter Exposure and Myocardial Ischemia in the Environmental Epidemiology of Arrhythmogenesis in the Women’s Health Initiative (EEAWHI) Study. *Environ Health Perspect*, 117(5), 751-756.
- <sup>xxix</sup> Pope AC, Burnett RT, Thun MJ, Calle EE, et al. (2002) Lung Cancer, cardiopulmonary mortality and long term exposure to fine particulate air pollution. *JAMA*, 287(9), 1132-1141
- <sup>xxx</sup> Miller AC, Arquilla B. (2008) Chronic diseases and natural hazards: impact of disasters on diabetic, renal and cardiac patients. *Prehosp Disaster Med*, 23(2).
- <sup>xxxi</sup> Chan EY, Sondorp E. (2007) Medical interventions following natural disasters: mission out on chronic medical needs. *Asia Pac J Public Health*, 19, 45-51.
- <sup>xxxii</sup> Wikipedia contributors. Bellevue Hospital Center. Wikipedia, The Free Encyclopedia. Available at: [http://en.wikipedia.org/wiki/Bellevue\\_Hospital\\_Center](http://en.wikipedia.org/wiki/Bellevue_Hospital_Center). Accessed August 14, 2014.
- <sup>xxxiii</sup> Uppal A, Evans L, Chitkara N, et al. (2013) In search of the silver lining: the impact of superstorm sandy on Bellevue hospital. *Ann AM Thorac Soc*, 10(2), 135-142.
- <sup>xxxiv</sup> Teperman S. (2013) Hurricane sandy and the greater New York health care system. *Journal of trauma and acute care surgery*, 74(6), 1401-1410.
- <sup>xxxv</sup> Curriero FC, Patz JA, Rose JB, Lele SD. (2001) The association between extreme precipitation and waterborne disease outbreaks in the United States, 1948–1994. *Am J Public Health*, 91 (8), 1194–1199.
- <sup>xxxvi</sup> Gaffield S, Goo R, Richards L, Jackson R. (2003) Public health effects of inadequately managed stormwater runoff. *American journal of public health*. 39 (9), 1527-33.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



<sup>xxxvii</sup> Umlauf, G., Bidoglio, G., Christoph, E., Kampheus, J., Krüger, F., Landmann, D., et al. (2005). The situation of PCDD/Fs and Dioxin-like PCBs after the flooding of river Elbe and Mulde in 2002. *Acta Hydrochimica et Hydrobiologica*, 33(5), 543–554.

<sup>xxxviii</sup> Manuel, J. (2006). In Katrina's wake. *Environmental Health Perspectives*, 114(1), 32–39.

<sup>xxxix</sup> Davis R. Hope turns to anguish at intensive-care unit. *USA Today*. 2005 September 16.

<sup>xl</sup> U.S. Energy Information Administration, Commercial Buildings Energy Consumption Survey 2012.

<sup>xli</sup> Meagan Brettelle, Health Canada Peter Berry, Health Canada Linda Varangu, Canadian Coalition for Green Health Care. PAHO, Health Care Facility Climate Change Resiliency Workshop Report. (2015) Page:6.

[http://www.paho.org/hq/index.php?option=com\\_docman&task=doc\\_view&Itemid=270&gid=32440&lang=en](http://www.paho.org/hq/index.php?option=com_docman&task=doc_view&Itemid=270&gid=32440&lang=en)

<sup>xlii</sup> A Report by the Government of the Commonwealth of Dominica. (2015) Rapid Damage and Impact Assessment Tropical Storm Erika. Page 12.

<https://www.gfdr.org/sites/default/files/publication/Commonwealth%20of%20Dominica%20-%20Rapid%20Damage%20and%20Needs%20Assessment%20Final%20Report%20.pdf>

<sup>xliii</sup> A Report by the Government of the Commonwealth of Dominica. (2015) Rapid Damage and Impact Assessment Tropical Storm Erika. Page 57.

<https://www.gfdr.org/sites/default/files/publication/Commonwealth%20of%20Dominica%20-%20Rapid%20Damage%20and%20Needs%20Assessment%20Final%20Report%20.pdf>

<sup>xliv</sup> Cox J and Abeku T. (2007) Early warning systems for malaria in Africa: from blueprint to practice. *Trends in parasitology*, 26(6), 243-246.

<sup>xlv</sup> Thomson MC, Dobkas-Reyes FJ, Mason SJ, Hagedorn R et al. (2006) Malaria early warnings based on seasonal climate forecasts from multi-model ensembles. *Nature*, 439, 576-579.

<sup>xlvi</sup> [No authors listed] Rift Valley Fever. World Health Organization Fact Sheet. Available at: <http://www.who.int/csr/disease/riftvalleyfev/en/>

<sup>xlvii</sup> Madani TA, Al-Mazrou YY, Al-Jeffri MH, Mishkhas AA, Al-Rabeah AM, Turkistani AM, Al-Sayed MO, Abodahish AA, Khan AS, Ksiazek TG, Shobokshi O. Rift Valley fever epidemic in Saudi Arabia: epidemiological, clinical, and laboratory characteristics. *Clinical Infectious Diseases*. 2003 Oct 15;37(8):1084-92.

<sup>xlviii</sup> Madani TA, Al-Mazrou YY, Al-Jeffri MH, Mishkhas AA, Al-Rabeah AM, Turkistani AM, Al-Sayed MO, Abodahish AA, Khan AS, Ksiazek TG, Shobokshi O. Rift Valley fever epidemic in Saudi Arabia: epidemiological, clinical, and laboratory characteristics. *Clinical Infectious Diseases*. 2003 Oct 15;37(8):1084-92.

<sup>xlix</sup> Rich KM, Wanyoike F. An assessment of the regional and national socio-economic impacts of the 2007 Rift Valley fever outbreak in Kenya. *The American journal of tropical medicine and hygiene*. 2010 Aug 5;83(2\_Suppl):52-7.

<sup>l</sup> [No authors listed] Rift Valley Fever Factsheet. Centers for Disease Control. Available at: <https://www.cdc.gov/vhf/rvf/index.html>

<sup>li</sup> Martin V, Chevalier V, Ceccato P, Anyamba A, De Simone L, Lubroth J, de La Rocque S, Domenech J. The impact of climate change on the epidemiology and control of Rift Valley fever. *Revue Scientifique et Technique-Office international des épizooties*. 2008;27(2):413-26.

<sup>lii</sup> Anyamba A, Linthicum KJ, Small J, Britch SC, Pak E, de La Rocque S, Formenty P, Hightower AW, Breiman RF, Chretien JP, Tucker CJ. Prediction, assessment of the Rift Valley fever activity in East and

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



---

Southern Africa 2006–2008 and possible vector control strategies. *The American journal of tropical medicine and hygiene*. 2010 Aug 5;83(2\_Suppl):43-51.

<sup>liii</sup> Linthicum KJ, Bailey CL, Davies FG, Tucker CJ. Detection of Rift Valley fever viral activity in Kenya, by satellite remote sensing imagery. *Science*. 1987 Mar 27;235:1656-60.

<sup>liv</sup> Pittiglio C, Pinto J, Munstermann S, Formenty P, del la Roche S, Raizman E, Beltran-Alcrudo D. El Nino and increasing risk of Rift Valley Fever – Warning to countries. *Empress Watch – Food and Agriculture Organization of the United Nations*. 2015; 34.

<sup>lv</sup> [No authors listed] Emerging Health Risk Notification. National Oceanic and Atmospheric Administration. 20 Dec 2015. Available at:  
<http://cpo.noaa.gov/sites/cpo/NOAA%20OneHealth/East%20Africa%20Rift%20Valley%20fever%20Risk%20notification%2020%20Dec%202015.pdf>

<sup>lvi</sup> Madani TA, Al-Mazrou YY, Al-Jeffri MH, Mishkhas AA, Al-Rabeah AM, Turkistani AM, Al-Sayed MO, Abodahish AA, Khan AS, Ksiazek TG, Shobokshi O. Rift Valley fever epidemic in Saudi Arabia: epidemiological, clinical, and laboratory characteristics. *Clinical Infectious Diseases*. 2003 Oct 15;37(8):1084-92.

<sup>lvi</sup> Utzinger J, Keiser J. (2006) Urbanization and tropical health – then and now. *Annals of Tropical Medicine and Parasitology*, 100(5), 517-533.

<sup>lviii</sup> Gottdenker N, Streicker D, Faust C, Carroll CR. (2014) Anthropogenic land use change and infectious diseases: A review of the evidence. *Ecohealth*.

<sup>lix</sup> Mutero CM, Kabutha C, Kimani V, Kabuage L, Gitau G, Ssenyonga J, Githure J, Muthami L, Kaida A, Musyoka L, Kiarie E, Oganda M. (2004) A transdisciplinary perspective on the links between malaria and agroecosystems in Kenya. *Acta Tropica*, 89(2), 171–186

<sup>lx</sup> Beldomenico P, Begon M. (2010) Disease spread, susceptibility and infection intensity. Vicious circles? *Trends in Ecology and Evolution*, 25(1), 21-27.

<sup>lxi</sup> Ryan ET. (2011) The cholera pandemic, still with us after half a century: Time to rethink. *PLoS Negl Trop*, 5(1).

<sup>lxii</sup> [No authors listed] (1998) Cholera outbreak among Rwandan refugees--Democratic Republic of Congo. *Centers for Disease Control, MMWR*, 47(19), 389-391.

<sup>lxiii</sup> Orata FD, Keimv PS, Boucher Y. (2014) The 2010 Cholera Outbreak in Haiti: How Science Solved a Controversy. *PLoS Pathog*, 10(4).

<sup>lxiv</sup> Ryan ET. (2011) The cholera pandemic, still with us after half a century: Time to rethink. *PLoS Negl Trop*, 5(1).

<sup>lxv</sup> Rose JB, Huffman, DE, Gennaccaro A. (2002) Risk and control of waterborne cryptosporidiosis. *Microbiology Reviews*, 26, 113-123

<sup>lxvi</sup> Goldstein ST, Juranek DD, Ravenholt O, Hightower AW, Martin DG, Mesnik JL, Griffiths SD, Bryant AJ, Reich RR, Herwaldt BL. (1996) Cryptosporidiosis: an outbreak associated with drinking water despite state-of-the-art water treatment, *Ann. Intern. Med*, 124(5), 459-68.

<sup>lxvii</sup> Adal KA, Sterling CR, Guerrant RL. *Cryptosporidium and related species*, in: *Infections of the Gastrointestinal Tract*, Raven Press, New York, 1995, pp. 1107–1128.

<sup>lxviii</sup> Dillingham RA, Lima AA, Guerrant, RL. (2002) Cryptosporidiosis: epidemiology and impact. *Microbes and infection*, (4), 1059-1066.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



- 
- <sup>lxxix</sup> Checkley W, Gilman RH, Epstein LD, Suarez M, Diaz JF, Cabrera L, Black RE, Sterling CR. (1997) Asymptomatic and symptomatic cryptosporidiosis: their acute effect on weight gain in Peruvian children. *Am. J. Epidemiol*, 145(2), 156–163.
- <sup>lxxx</sup> Dillingham RA, Lima AA, Guerrant, RL. (2002) Cryptosporidiosis: epidemiology and impact. *Microbes and infection*, (4) 1059-1066.
- <sup>lxxxi</sup> MacKenzie WR, Hoxie NJ, Proctor ME, Gradus S, Blair KA, Peterson DE, Kazmierczak JJ, Addiss DG, Fox KR, Rose JB, David JP. (1994) A massive outbreak in Milwaukee of *Cryptosporidium* infection transmitted through the public water supply. *New Engl. J. Med*, 331(3), 161-167.
- <sup>lxxxii</sup> [No authors listed] (2012) Progress on drinking water and sanitation: UNICEF/WHO Joint Monitoring Programme update.
- <sup>lxxxiii</sup> Hoagland P, Lara DI, Polansky LY, Kerkpatrick et al. The costs of respiratory illnesses arising from Florida gulf coast *Karenia brevis* blooms. *Environmental Health Perspectives*. 2009. Aug Vol 117 No8.
- <sup>lxxxiv</sup> Heil CA, Steidinger KA. Monitoring, management and mitigation of *Kareina* blooms in the eastern Gulf of Mexico. *Harmful Algae*. 2009 8:611–617.
- <sup>lxxxv</sup> Fleming LE, Kirkpatrick B, Backer LC, Bean JA, Wanner A, Dalpra D, et al. Initial evaluation of the effects of aerosolized Florida red tide toxins (brevetoxins) in persons with asthma. *Environ Health Perspect* 2005 113:650–657.
- <sup>lxxxvi</sup> Kirkpatrick B, Fleming LE, Backer LC, Bean JA, Tamer R, Kirkpatrick G, et al. Environmental exposures to Florida red tides: effects on emergency room respiratory diagnoses admissions. *Harmful Algae*. 2006. 5:526–533.
- <sup>lxxxvii</sup> Watts N, Amann M, Ayeb-Karlsson S, Belesova K, Bouley T, Boykoff M, Byass P, Cai W, Campbell-Lendrum D, Chambers J, Cox PM. The Lancet Countdown on health and climate change: from 25 years of inaction to a global transformation for public health. *The Lancet*. 2017 Oct 30.
- <sup>lxxxviii</sup> Food and Agricultural Organization of the United Nations. The state of food insecurity in the world: Food insecurity – when people must live with hunger and fear of starvation. Rome, Food and Agriculture Organization, 2002.
- <sup>lxxxix</sup> Pitt M, Khandker SR, Chowdhury OH, Millimet D. (2003) Credit programs for the poor and the health status of children in rural Bangladesh. *International economic review*, 44(1).
- <sup>lxxx</sup> Rahman A. Women and microcredit in rural Bangladesh: An anthropological study of Grameen Bank lending. Westview Press; 2008 Dec 15.
- <sup>lxxxii</sup> Warren K, Bundy DAP, Anderson R, Davis AR, Henderson DA, Jamison DT, Prescott N, Senft A. (1993) Helminth Infection. Jamison DT, Mosley W, Measham A, Bobadilla J, editors. Disease control priorities in developing countries. Oxford: Oxford Medical Publications.
- <sup>lxxxiii</sup> Fernando MA, Balasuriya S. (1983) Effect of *Ascaris lumbricoides* infestation on growth of children. *Indian Pediatrics*, 20(10), 721–31.
- <sup>lxxxiiii</sup> Stephenson L. (1987) The impact of helminth infections on human nutrition. London: Taylor and Francis.
- <sup>lxxxv</sup> Nokes C and Bundy DA. (1994) Does helminth infection affect mental processing and education achievement? *Parasitology today*, 1(10), 14-18.
- <sup>lxxxvi</sup> Cunningham-Rundles S, McNeeley DF, Moon A. (2005) Mechanisms of nutrient modulation of the immune response. *J Allergy Clin Immunol*, 115, 119-28.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



---

<sup>lxxxvi</sup> Brito LL, Barreto ML, Silva Rde C, et al. (2006) Moderate- and low-intensity co-infections by intestinal helminths and *Schistosoma mansoni*, dietary iron intake, and anemia in Brazilian children. *Am J Trop Med Hyg*, 75, 939-44.

<sup>lxxxvii</sup> Papier K, Williams GM, Luceres-Catubig R, Ahmed F, Olveda R et al. (2014) Childhood malnutrition and parasitic helminth interactions. *Clinical infectious disease*, 59(2), 234-243.

<sup>lxxxviii</sup> [No authors listed] (2012) Pesticide Exposure in Children. Council on Environmental Health. *Pediatrics*, 130, 2012-2757.

<sup>lxxxix</sup> Kimmel CA, Collman GW, Fields N, Eskenazi B. (2005) Lessons learned for the National Children's Study from the National Institute of Environmental Health Sciences/U.S. Environmental Protection Agency Centers for Children's Environmental Health and Disease Prevention Research. *Environ Health Perspect*, 113(10), 1414-1418.

<sup>xc</sup> Eckerman DA, Gimenes LA, de Souza RC, Lopes Galvão PR, Sarcinelli PN, Chrisman JR. (2007) Age related effects of pesticide exposure on neurobehavioral performance of adolescent farm workers in Brazil. *Neurotoxicol Teratol*, 29(1), 164-175.

<sup>xc1</sup> Landrigan PJ, Claudio L, Markowitz SB, et al. (1999) Pesticides and inner-city children: exposures, risks, and prevention. *Environ Health Perspect*, 107(3), 431-437

<sup>xcii</sup> Perera F. (2014) Children's environmental health: A critical challenge of our time. *Lancet*, 383(9921), 943-944.

<sup>xciii</sup> Perera F. (2014) Children's environmental health: A critical challenge of our time. *Lancet*, 383(9921), 943-944..

<sup>xciv</sup> Batisti DS, Naylor RL. (2009) Historical warnings of future food insecurity with unprecedented seasonal heat. *Science*, 323 (5911), 240-244.

<sup>xcv</sup> Bryce J, Coitinho D, Darnton-Hill, I, Pelletier D, Pinstrup-Andersen P. (2008) Maternal and child undernutrition: effective action at national level. *Lancet*, 371(9611) 510-26.

<sup>xcvi</sup> Bryce J, Coitinho D, Darnton-Hill, I, Pelletier D, Pinstrup-Andersen P. (2008) Maternal and child undernutrition: effective action at national level. *Lancet*, 371(9611) 510-26.

<sup>xcvii</sup> Black, RE, Allen, LH, Bhutta ZA, Caulfield LE, et al. (2008) Maternal and child undernutrition: global and regional exposures and health consequences. *Lancet*, 371(9608), 234-60.

<sup>xcviii</sup> Hoddinott J, Behrman JR, Maluccio JA, et al. (2008) Effect of a nutrition intervention during early childhood on economic productivity in Guatemalan adults. *Lancet*, 371(9610), 411-416.

<sup>xcix</sup> Nelson Textbook of Pediatrics, 19<sup>th</sup> edition. 2011. Nutrition, Food Security and Health.

<sup>c</sup> [No authors listed] Marine Toxins. National Center or Emerging and Zoonotic Infectious Diseases. Centers for Disease Control. 2013. Accessed July 17, 2013.

[http://www.cdc.gov/nczved/divisions/dfbmd/diseases/marine\\_toxins/#diseases](http://www.cdc.gov/nczved/divisions/dfbmd/diseases/marine_toxins/#diseases)

<sup>ci</sup> Lehane L, Lewis RJ. Ciguatera: recent advances but the risk remains. *Int J Food Microbiol*. 2000; 61:91.

<sup>cii</sup> Lewis RJ, Sellin M, Poli MA, et al. Purification and characterization of ciguatoxins from moray eel (*Lycodontis javanicus*, Muraenidae). *Toxicon*. 1991; 29:1115-1127.

<sup>ciii</sup> Tunik M. Goldfrank's Toxicologic Emergencies, 9e. McGraw Hill Companies. 2011.

<sup>civ</sup> Doak CM, Adair LS, Bentley M, Monteiro C, Popkin BM. (2005) The dual burden household and the nutrition transition paradox. *International Journal of Obesity*, 29, 129-136.

## Educational Materials

A Clinical Health Care Student Exploration of the  
Global Impacts of Climate Change on Human Health



- 
- <sup>cv</sup> Popkin BM, Adair L, Ng SW. (2012) Now and then: the global nutrition transition: the pandemic obesity in developing countries. *Nutrition Review*, 70(1), 3-21.
- <sup>cvi</sup> Popkin BM. (1999) Urbanization, Lifestyle Changes and the Nutrition Transition. *World Development*, 27, 1905-1916.
- <sup>cvi</sup> Popkin, BM, and Doak CM. (1998) The Obesity Epidemic Is a Worldwide Phenomenon. *Nutrition Reviews*, 56, 106-114.
- <sup>cvi</sup> Kelly T, Yang W, Chen CS, Reynolds K, He J. (2008) Global Burden of obesity in 2005 and projections to 2030. *International Journal of Obesity*, 32(9), 1431-7.
- <sup>cix</sup> [No authors listed] (2011) Human resources for mental health: workforce shortages in low and middle-income countries. *World health organization, Human Resources for Health Observer*, 8.
- <sup>cx</sup> Felder SS, Seligman J, Burrows-McElwain CK, Robinson ME, Hierholzer E. (2014) Disaster Trauma: Federal resources that help communities on their road to recovery. *Disaster Med Public Health Prep*, 1(5).
- <sup>cx</sup> Chan CS, Rhodes JE. (2014) Measuring exposure in Hurricane Katrina: a meta-analysis and an integrative data analysis. *PLoS One*, 9(4).
- <sup>cxii</sup> Keenan HT, Marshall SW, Nocera MA, et al. (2004) Increased incidence of inflicted traumatic brain injury in children after a natural disaster. *Am J Prev Med*, 26,189–93.
- <sup>cxiii</sup> Krug EG, Kresnow M, Peddicord JP, et al. (1998) Suicide after natural disasters. *N Engl J Med*, 338, 373-8.
- <sup>cxiv</sup> Prince M, Patel V, Saxena S, Maj M, Maselko J, Phillips M, Rahman A. (2007) No health without mental health. *Lancet*, 370 (9590), 859-877.
- <sup>cxv</sup> Thomas K, Ellis A, Konrad T, Holzer C, Morrissey J. (2009) County-level estimates of mental health professional shortage in the United States. *Psychiatric Services*, 60 (10), 1323-8.
- <sup>cxvi</sup> Mollica R. (2004) Surviving Torture. *New England Journal of Medicine*, 351(5-7).
- <sup>cxvii</sup> Kip K, Rosenzweig L, Hernandez D, et al. (2013) Accelerated resolution therapy for treatment of pain secondary to symptoms of combat-related posttraumatic stress disorder. *European Journal of Psychotraumatology*, 5, 240-66.
- <sup>cxviii</sup> Plagge J, Lu MW, Lovejoy TI, Karl AI, Dobscha S. (2013) Treatment of comorbid pain and PTSD in returning veterans: a collaborative approach utilizing behavioral activation. *Pain Medicine*, 14, 1164-1172.
- <sup>cxix</sup> Haushofer J and Fehr E. (2014) On the psychology of poverty. *Science*, 344(6186), 862-867.
- <sup>cxx</sup> Filipovsky J, Ducimetiere P, Eschwege E, Richard JL, Rosselin G, Claude JR. (1996) The relationship of blood pressure with glucose, insulin, heart rate, free fatty acids and plasma cortisol levels according to degree of obesity in middle-aged men. *Journal of hypertension*, 14(2), 229-35.
- <sup>cxxi</sup> Damti OB, Sarid O, Sheiner E, Zilberstein T, Cwikel J. (2008) Stress and distress in infertility among women. *Harefuah*, 147(3), 256-60.
- <sup>cxxii</sup> Laat, C and Haushofer J. (2013) Poverty and stress: Rainfall shocks increase levels of the stress hormone cortisol. *Massachusetts Institute of Technology Working Paper*.
- <sup>cxxiii</sup> Arnetz BB, Brenner SO, Levi L, Hjelm R, et al. (1991) Neuroendocrine and immunologic effects of unemployment and job insecurity. *Psychother psychosom*, 55, 76–80.

## Educational Materials

*A Clinical Health Care Student Exploration of the Global Impacts of Climate Change on Human Health*



---

<sup>cxiv</sup> Nabi H, Kivimaki M, Batty GD, Shipley MJ, Britton A, Brunner EJ, Cahtera J, Lemogne C, Elbaz A, Singh-Manoux A. (2013) Increased risk of coronary heart disease among individuals reporting adverse impact of stress on their health: the Whitehall II prospective cohort study. *European Heart Journal*, 34(34), 2697-705.

<sup>cxv</sup> Rodney N, Mullican C. (2014) A biocultural study of the effects of maternal stress on mother and newborn health in the Democratic Republic of Congo. *American Journal of Physical Anthropology*, 155(2), 200-209.

<sup>cxvi</sup> Litz, B. (2004) *Early intervention for trauma and traumatic loss*. New York: Guilford Press.

<sup>cxvii</sup> Tedeschi RG, Calhoun LG. (1996) The Posttraumatic Growth Inventory: Measuring the positive legacy of trauma. *Journal of Traumatic Stress*, 9, 455–471.

<sup>cxviii</sup> Walsh F. (2007) *Traumatic Loss and Major Disasters: Strengthening Family and Community Resilience*. *Family Process*, 46(2).

<sup>cxix</sup> Landau J, Saul J. (2004) Family and community resilience in response to major disaster. In *Living beyond loss: Death in the family* (2nd ed) F. Walsh & M. McGoldrick (Eds.) New York, Norton. pp. 285–309.

<sup>cxx</sup> United States Department of Health and Human Services. *Sustainable and Climate Resilient Healthcare Facilities Initiative*. Available at: <https://toolkit.climate.gov/tool/sustainable-and-climate-resilient-health-care-facilities-toolkit>. Accessed: 12/3/17