Detection of Survivin (BIRC5) in Formalin-Fixed, Paraffin-Embedded Mouse Tissue

Reagent and Antibody Information

1X Wash Buffer
3% Hydrogen Peroxide
1% BSA Diluent
1X Citrate Buffer
Normal Rabbit IgG – Affinity Purified
DAB Chromogen
Hematoxylin

Blocking Serum: Normal Donkey Serum
Jackson Immunoresearch Laboratories, Inc.
West Grove, PA 19390
www.jacksonimmuno.com
1-800-367-5296
Catalog # 017-000-011

Avidin / Biotin Blocking Kit
Vector Laboratories, Inc.
Burlingame, CA 94010
www.vectorlabs.com
1-800-227-6666
Catalog # SP-2001

Primary Antibody: Rabbit Polyclonal Anti-Survivin Antibody
Abcam, Inc
Cambridge, MA 02139
www.abcam.com
1-888-772-2226
Catalog # ab469

Secondary Antibody: Donkey Anti-Rabbit IgG (H+L) Biotin-SP-Conjugated
Jackson Immunoresearch Laboratories, Inc.
West Grove, PA 19390
www.jacksonimmuno.com
1-800-367-5296
Catalog # 711-065-152

Label Complex: R.T.U. Vectastain Elite ABC Reagent
Vector Laboratories, Inc.
Burlingame, CA 94010
www.vectorlabs.com
1-800-227-6666
Catalog # PK-7100

Staining Procedure

Positive Control Tissue: Liver
Stain Localization: Cytoplasmic – peri-venular

1. Deparaffinize and hydrate slides through the following solutions:

Solution	Repetitions	Time
Xylene	2 times	5 minutes
100% Ethanol	2 times	3 minutes
95% Ethanol	2 times	3 minutes
1X Wash Buffer	2 times	5 minutes

2. Quench endogenous peroxidase by placing the slides in 3% hydrogen peroxide for 15 minutes.

3. Rinse slides in 2 changes of 1X wash buffer for 5 minutes each.

4. Heat-Induced Epitope Retrieval Using The Decloaker
   Add 500 ml of distilled water to the pan inside the decloaker.
   Place a full rack of slides into a Tissue Tek® container with 200 ml of 1X citrate buffer
   (Insert blank slides into any empty slots in the rack to ensure even heating of slides)
   Place the container stably inside the pan and decloak for 5 minutes. *Maximum Pressure* _____
   Depressurize for 10 minutes.
   Remove pan top and cool for 10 minutes. *Temperature Before Cooling Slides*_____
   Rinse the slides in 2 changes of distilled water for 3 minutes each time.

5. Rinse slides in 2 changes of 1X wash buffer for 5 minutes each.

6. Block with 10% normal donkey serum for 20 minutes at room temperature.
   Lot #_____ Date Reconstituted_____

   DO NOT RINSE SLIDES. CONTINUE TO AVIDIN-BIOTIN BLOCK.

7. Avidin / Biotin Blocking Kit
   Lot #_____ Exp. Date_____ New Kit: yes / no
   Apply avidin block for 15 minutes at room temperature.
   Quick rinse in 1X wash buffer.
   Apply biotin block for 15 minutes at room temperature.

   DO NOT RINSE SECTIONS WITH BUFFER BEFORE ADDING PRIMARY ANTIBODY.
   ONLY WIPE EXCESS BLOCK.

8. Apply primary antibody at a 1:500 dilution. Incubate for 1 hour at room temperature.
   Lot #_____ Exp. Date _____

   For negative control slides, dilute normal rabbit IgG so that it's IgG protein concentration matches that of the primary antibody (if necessary). Then make a 1:500 dilution. If the concentrations can't be matched using this method, the dilution for the negative reagent may need to be adjusted. Apply the

negative and incubate for 1 hour at room temperature.
Lot #_____ Exp. Date _____

9. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

10. Apply the donkey anti-rabbit secondary antibody at a 1:500 dilution. Incubate for 30 minutes at room temperature.
    Lot #_____ Date Reconstituted _____

11. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

12. Apply the Vectastain R.T.U. Elite Label and incubate for 30 minutes at room temperature.
    Exp. Date_____ New Kit: yes / no

13. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each time.

14. Apply the DAB chromogen. Incubate in the dark for 6 minutes at room temperature.
    (Add 1 drop of DAB per ml of substrate)
    Lot #_____ Exp. Date_____ New Kit: yes / no

15. Rinse the slides in tap water 3 minutes.

16. Counterstain with hematoxylin for 20 seconds.

17. Rinse the slides in tap water until water is clear.

18. Gently agitate slides in 1X wash buffer until the tissues turn blue.

19. Dehydrate through the following solutions:

Solutions	Repetitions	Time
95% Ethanol	1 time	3 minutes
100% Ethanol	3 times	3 minutes
Xylene	2 times	5 minutes

20. Coverslip

Updated 08/23/10

This protocol has been made available for use by others. We cannot guarantee optimal results, so staining conditions must be determined by the end user.
Protocol modifications may be required due to lot number changes and/or reagent substitutions.