

Concept Clearance

Branch: Population Health Branch

Council Period: June 2019

Concept Title: "Research to Action: Assessing and Addressing Community Exposures to Environmental Contaminants"

Introduction

We are proposing the renewal of the Research to Action (R2A) program. The R2A program promotes and advances academic-community partnerships to address environmental health concerns of diverse communities and to ensure authentic community participation in research leading to public health actions. The program has evolved from an R21 solicitation (RFA, 2008) to two iterations of an R01 solicitation (PA, 2012 and 2016) that requires the environmental health problem and development of research questions be informed by community members or organizations. All projects must demonstrate an established community-university collaboration, utilize multidisciplinary teams to address the many interacting factors contributing to environmental health risks, and disseminate findings and conduct public health actions that are culturally appropriate for the communities and subpopulations involved in the research.

Background information: Environmental threats such as the rapidly increasing number of fracking operations in the US and the numerous communities who have discovered lead, PFAS and other types of chemical or metal contamination in local water sources, suggest that continued and well-publicized exposure to toxic substances will remain a major public health issue (Denham et al, 2019; Rappazzo et al, 2017; Boronow et al. 2019). This will be true not only for those at the lowest socioeconomic status and for specific ethnic and racial subpopulations but for entire communities or regions. GenX and other chemicals released into local North Carolina rivers, for example represents an indiscriminate exposure that has affected many diverse communities and populations in the State. The media attention to this environmental disaster and to other natural and manmade disasters in the US has greatly increased public interest in environmental health research which the Research to Action program is poised to help address.

However, environmental risks continue to disproportionately affect certain ethnic and racial subpopulations. These environmental health disparities, which cannot be fully explained by socio-cultural or behavioral factors, persist throughout the United States. In Ohio, for example, a recent study showed that Class II wells that receive fracking wastewater are disproportionately situated in lower-income and rural communities, areas with the greatest social vulnerability (Kunkel, 2019). And in areas such as Appalachia, public health studies have focused primarily on the known impacts of coal mining to the general population but have not fully explored the contributions of socioeconomic status, built environment, and food deserts, personal exposure, and other sources of air, water and soil pollution, thus limiting the ability to identify causal relationships between environmental and social conditions and public health (Krometis et al., 2017). Research in air pollution worldwide has likewise revealed the disproportionate effects on certain subgroups defined by sociodemographics (race/ethnicity, age, sex and socioeconomic status [SES]), geographic location (rural vs. urban), comorbid conditions and economic conditions, all factors which contribute to the adverse cardiovascular effects of exposure to ambient air pollution (Tibuakuu et al., 2018).

The Research to Action program is therefore targeted to addressing not just environmental threats but environmental health disparities. Since the 1990s, numerous studies have demonstrated the ongoing health disparities experienced by African-Americans, particularly those living in inner cities – this awareness of disparities was coupled with a growing awareness of the role of social determinants of health (SDOH). And recent studies by NIEHS funded investigators are confirming that environmental health disparities in exposures to drinking water contaminants (e.g. nitrates) are related to race and ethnicity (Schneider et al., 2019) or to air pollution exposures from the built environment, goods movements or proximity to major roadways (Burwell et al, 2017; Wilson et al, 2015). Although genetics can account for some of the disparities (Daya and Barnes, 2019), other known causes of these disparities include multiple and chronic environmental exposures, and SDOHs e.g., socioeconomic status, the built environment, population density, noise and violence and access to nutritious affordable food. Many low SES populations experience these disparities however, studies have also shown that racial discrimination, and even perceived racial discrimination, is a potent stressor that has been linked to psychosocial stress and poor physical health (Baraja et al, 2019), adding yet another factor to heightened risks for disease among African-Americans. Several of the R2A projects have been addressing the multiple exposures and social conditions that inner city African-Americans (AA) experience over time. In one of these studies, the effects of unregulated metal recycling on air quality in low SES neighborhoods in Houston is being explored. In another R2A study the PI has coupled with local AA community leaders, academic investigators and public health practice partners in research that is culminating in the development and implementation of a public health action plan that reduces exposure to air pollutants, specifically sources of PM2.5 and mitigates the adverse health effects of such exposures.

Another particularly vulnerable subpopulation are immigrants. Environmental health disparities remain high among Latinx (a gender-neutral version of Latino or Latina) groups migrating to the US to work in farming, industry and domestic cleaning jobs as these occupations are often the most harmful in terms of environmental exposures. Research to Action projects and other NIEHS funded research among farmworkers has shown the consistently negative impacts of pesticide exposure among workers (Stein et al., 2016; Arcury et al., 2018) as well as their family members (Vidi et al., 2017). And, a recently funded R2A study is exploring ways to mitigate the multiple chemical exposures domestic cleaners typically experience in their jobs.

Although, many ethnic and racial subpopulations in the US experience disproportionate exposures to environmental risks, Native Americans remain among the most vulnerable of populations (Abbasi, 2018; Kunitz et al., 2014). Twenty-five percent of Superfund sites have been identified on tribal lands; climate change is rapidly damaging indigenous inhabited ecosystems in the Arctic, Gulf of Mexico and American Southwest among other regions; and increased resource extraction in proximity to tribal communities continues to expose these populations to ongoing and multiple exposures that threaten health (Byrne et al., 2018; McOliver et al, 2015). Native Americans living near abandoned uranium mines, for example, have an increased likelihood for kidney disease and hypertension, and for developing multiple chronic diseases linked to their proximity to the mine waste (Lewis et al., 2017).

The Research to Action program, with its requirement to engage with local community members and organizations, is thus uniquely poised to not only address environmental exposures affecting health, but to doing so in a culturally appropriate manner for specific racial/ethnic subpopulations, as well as for low SES, rural remote or inner-city communities.

We believe that the R2A program exemplifies the importance of community involvement in environmental health research and will continue to reach the most vulnerable subpopulations and address some of the more severe environmental health disparities in the US. The program additionally represents a validated approach for sustaining public health actions in vulnerable communities and for raising the environmental health literacy of affected community members (as well as stimulating interest in students in STEM topics).

Bibliography

Denham A, Willis M, Zavez A, Hill E. Unconventional natural gas development and hospitalizations: evidence from Pennsylvania, United States, 2003-2014. *Public Health*. 2019 Mar;168:17-25. doi: 10.1016/j.puhe.2018.11.020. Epub 2019 Jan 21. PubMed PMID: 30677623; PubMed Central PMCID: PMC6408296.

Rappazzo KM, Coffman E, Hines EP. Exposure to Perfluorinated Alkyl Substances and Health Outcomes in Children: A Systematic Review of the Epidemiologic Literature. *Int J Environ Res Public Health*. 2017 Jun 27;14(7). pii: E691. doi: 10.3390/ijerph14070691. Review. PubMed PMID: 28654008; PubMed Central PMCID: PMC5551129.

Boronow KE, Brody JG, Schaidler LA, Peaslee GF, Havas L, Cohn BA. Serum concentrations of PFASs and exposure-related behaviors in African American and non-Hispanic white women. *J Expo Sci Environ Epidemiol*. 2019 Mar;29(2):206-217. doi: 10.1038/s41370-018-0109-y. Epub 2019 Jan 8. PubMed PMID: 30622332; PubMed Central PMCID: PMC6380931.

Konkel L. Where Does Fracking Wastewater Go? Socioeconomic Predictors of Class II Well Placement in Ohio. *Environ Health Perspect*. 2019 Mar;127(3):34003. doi: 10.1289/EHP4797. PubMed PMID: 30912995.

Krometis LA, Gohlke J, Kolivras K, Satterwhite E, Marmagas SW, Marr LC. Environmental health disparities in the Central Appalachian region of the United States. *Rev Environ Health*. 2017 Sep 26;32(3):253-266. doi: 10.1515/reveh-2017-0012. Review. PMID: 28682789.

Tibuakuu M, Michos ED, Navas-Acien A, Jones MR. Air Pollution and Cardiovascular Disease: A Focus on Vulnerable Populations Worldwide. *Curr Epidemiol Rep*. 2018 Dec;5(4):370-378. doi: 10.1007/s40471-018-0166-8. Epub 2018 Sep 19. PubMed PMID: 30931239; PubMed Central PMCID: PMC6435302.

Daya M, Barnes KC. African American ancestry contribution to asthma and atopic dermatitis. *Ann Allergy Asthma Immunol*. 2019 Feb 15. pii: S1081-1206(19)30101-2. doi: 10.1016/j.anai.2019.02.009. [Epub ahead of print] Review. PubMed PMID: 30772392.

Schaidler LA, Swetschinski L, Campbell C, Rudel RA. Environmental justice and drinking water quality: are there socioeconomic disparities in nitrate levels in U.S. drinking water? *Environ Health*. 2019 Jan 17;18(1):3. doi: 10.1186/s12940-018-0442-6. PubMed PMID: 30651108; PubMed Central PMCID: PMC6335696.

Burwell-Naney K, Wilson SM, Tarver SL, Svendsen E, Jiang C, Ogunsakin OA, Zhang H, Campbell D, Fraser-Rahim H. Baseline Air Quality Assessment of Goods Movement Activities before the Port of Charleston Expansion: A Community-University Collaborative. *Environ Justice*. 2017 Feb 1;10(1):1-10. doi: 10.1089/env.2016.0018. PubMed PMID: 29576842; PubMed Central PMCID: PMC5363220.

Wilson S, Burwell-Naney K, Jiang C, Zhang H, Samantapudi A, Murray R, Dalemarre L, Rice L, Williams E. Assessment of sociodemographic and geographic disparities in cancer risk from air toxics in South Carolina. *Environ Res*. 2015 Jul;140:562-8. doi: 10.1016/j.envres.2015.05.016. Epub 2015 May 28. PubMed PMID: 26037107; PubMed Central PMCID: PMC4492882.

Barajas CB, Jones SCT, Milam AJ, Thorpe RJ Jr, Gaskin DJ, LaVeist TA, Furr-Holden CDM. Coping, Discrimination, and Physical Health Conditions Among Predominantly Poor, Urban African Americans: Implications for Community-Level Health Services. *J Community Health*. 2019 Mar 26. doi: 10.1007/s10900-019-00650-9. [Epub ahead of print] PubMed PMID: 30915675.

Abbasi, J. *JAMA*. 2018 Jan 9;319(2):109-111. doi: 10.1001/jama.2017.10122. Why Are American Indians Dying Young? DOI: 10.1001/jama.2017.10122 PMID: 29318275 [Indexed for MEDLINE]

Kunitz SJ, Veazie M, Henderson JA. Historical trends and regional differences in all-cause and amenable mortality among American Indians and Alaska Natives since 1950. *Am J Public Health*. 2014 Jun;104 Suppl 3:S268-77. doi: 10.2105/AJPH.2013.301684. Epub 2014 Apr 22. PubMed PMID: 24754651; PubMed Central PMCID: PMC4035869.

Byrne SC, Miller P, Seguinot-Medina S, Waghiyi V, Buck CL, von Hippel FA, Carpenter DO. Associations between serum polybrominated diphenyl ethers and thyroid hormones in a cross-sectional study of a remote Alaska Native population. *Sci Rep*. 2018 Feb 2;8(1):2198. doi: 10.1038/s41598-018-20443-9. PubMed PMID: 29396447; PubMed Central PMCID: PMC5797183.

McOliver CA, Camper AK, Doyle JT, Eggers MJ, Ford TE, Lila MA, Berner J, Campbell L, Donatuto J. Community-based research as a mechanism to reduce environmental health disparities in american Indian and alaska native communities. *Int J Environ Res Public Health*. 2015 Apr 13;12(4):4076-100. doi: 10.3390/ijerph120404076. PubMed PMID: 25872019; PubMed Central PMCID: PMC4410234.

Lewis J, Hoover J, MacKenzie D. Mining and Environmental Health Disparities in Native American Communities. *Curr Environ Health Rep*. 2017 Jun;4(2):130-141. doi: 10.1007/s40572-017-0140-5. Review. PMID: 28447316; PubMed Central PMCID: PMC5429369.

Stein LJ, Gunier RB, Harley K, Kogut K, Bradman A, Eskenazi B. Early childhood adversity potentiates the adverse association between prenatal organophosphate pesticide exposure and child IQ: The CHAMACOS cohort. *Neurotoxicology*. 2016 Sep;56:180-187. doi: 10.1016/j.neuro.2016.07.010. Epub 2016 Jul 26. PubMed PMID: 27474229; PubMed Central PMCID: PMC5765748.

Cushing L, Faust J, August LM, Cendak R, Wieland W, Alexeeff G. Racial/Ethnic Disparities in Cumulative Environmental Health Impacts in California: Evidence From a Statewide Environmental Justice Screening Tool (CalEnviroScreen 1.1). *Am J Public Health*. 2015 Nov;105(11):2341-8. Doi 10.2105/ AJPH.2015.302643. Epub 2015 Sep 17. PMID: 26378826; PMCID: PMC4605180.

Arcury TA, Laurienti PJ, Talton JW, Chen H, Howard TD, Barr DB, Mora DC, Quandt SA. Pesticide Urinary Metabolites Among Latina Farmworkers and Nonfarmworkers in North Carolina. *J Occup Environ Med*. 2018 Jan;60(1):e63-e71. doi: 10.1097/JOM.0000000000001189. PubMed PMID: 29023343; PubMed Central PMCID: PMC5758422.

Vidi PA, Anderson KA, Chen H, Anderson R, Salvador-Moreno N, Mora DC, Poutasse C, Laurienti PJ, Daniel SS, Arcury TA. Personal samplers of bioavailable pesticides integrated with a hair follicle assay of DNA damage to assess environmental exposures and their associated risks in children. *Mutat Res*. 2017 Oct;822:27-33. doi: 10.1016/j.mrgentox.2017.07.003. Epub 2017 Jul 16. PubMed PMID: 28844239; PubMed Central PMCID: PMC5607735.

Research Goals and Scope

The research goals and scope remain the same as the previous Program Announcements.

The two main objectives are:

- 1) to conduct research to collect and characterize information about environmental health concerns of significance to a community and
- 2) to develop and implement a strategy to translate and disseminate research findings to community members, public health professionals and/or policymakers to support an action that will ultimately promote the reduction of exposure and reduce the health impact from environmental stressors.

To meet these stated objectives, applicants should propose Community Engaged Research (CErR) projects that incorporate three elements - research, community engagement and public health action. Evaluation is an optional element.

Mechanism and Justification

Due to recent changes in NIH policy about Program Announcements, we are requesting consideration of an annual Request for Applications (RFA) for the Research to Action (R2A) program.

Through an annual receipt date, the program will continue to provide flexibility to respond to emerging concerns and may also stimulate interest in new communities to build capacity to address their local concerns

We are proposing an R01 mechanism for this RFA and a timeline as follows:

Jun 2019	Present program concept to NAEHS Council
Sep 2019	Publish FOAs
Nov 2019	Receipt of applications
Spring 2020	Reviews
Sep 2020	Present program plan to NAEHS Council
Late Fall (FY2021)	Awards

- **Mechanism:** R01
- **FOA:** RFA issued for 3 years
- **Anticipated number of awards:** 5-6
- **Total cost for program:**
 - \$500,000 per award for 5 years
 - \$3 million total annually