

Detection of Cytokeratin 5/8 in Formalin-Fixed, Paraffin-Embedded Rat Tissue

Reagent and Antibody Information

[1X Wash Buffer](#)

[3% Hydrogen Peroxide](#)

[1% BSA Diluent](#)

[Trypsin](#)

[DAB Chromogen](#)

[Hematoxylin](#)

Blocking Serum: Normal Horse Serum

Jackson ImmunoResearch Laboratories, Inc.

West Grove, PA 19390

www.jacksonimmuno.com

1-800-367-5296

Catalog # 008-000-001

Avidin / Biotin Blocking Kit

Vector Laboratories, Inc.

Burlingame, CA 94010

www.vectorlabs.com

1-800-227-6666

Catalog # SP-2001

Primary Antibody: Mouse Monoclonal Cytokeratin 5/8 Antibody (RCK102)

Santa Cruz Biotechnology, Inc.

Santa Cruz, CA 95060

www.scbt.com

1-800-457-3801

Catalog # sc-32328

Negative Control Serum: Purified Mouse IgG1 Isotype Control Serum

BD Biosciences

San Jose, CA 95131

www.bdbiosciences.com

1-855-236-2772

Catalog # 557273

Secondary Antibody: Biotinylated Horse Anti-Mouse IgG (H+L)

Vector Laboratories, Inc.

Burlingame, CA 94010

www.vectorlabs.com

1-800-227-6666

Catalog # BA-2001

Label Complex: R.T.U. Vectastain Elite ABC Reagent
Vector Laboratories, Inc.
Burlingame, CA 94010
www.vectorlabs.com
1-800-227-6666
Catalog # PK-7100

Staining Procedure

Positive Control Tissue: Female reproductive tract – glandular epithelium of the uterus and cervix
Stain Localization: Cytoplasmic

1. Deparaffinize and hydrate slides through the following solutions:

Solution	Repetitions	Time
Xylene	2 times	5 minutes
100% Ethanol	2 times	3 minutes
95% Ethanol	2 times	3 minutes
1X Wash Buffer	2 times	5 minutes

2. Quench endogenous peroxidase by placing the slides in 3% hydrogen peroxide for 15 minutes.

3. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

4. Proteolytic-Induced Epitope Retrieval Using Trypsin

Incubate the slides in a **0.01% trypsin** solution in a water bath at 37°C for 20 minutes.

(DO NOT add the trypsin to the 0.05M Tris-HCl • CaCl₂ solution until 5 minutes prior to incubation.

Trypsin loses 75% of its reactivity within 30 minutes at 37°C.)

Rinse the slides in distilled water for 1 minute to stop the enzymatic digestion.

5. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

6. Block with 10% normal horse serum for 20 minutes at room temperature.

Lot # _____ Date Reconstituted _____

DO NOT RINSE SLIDES. CONTINUE TO AVIDIN-BIOTIN BLOCK.

7. Avidin / Biotin Blocking Kit

Lot # _____ Exp. Date _____ New Kit: yes / no

Apply avidin block for 15 minutes at room temperature.

Quick rinse in 1X wash buffer.

Apply biotin block for 15 minutes at room temperature.

DO NOT RINSE SECTIONS WITH BUFFER BEFORE ADDING PRIMARY ANTIBODY.
ONLY WIPE EXCESS BLOCK.

8. Apply primary antibody at a 1:25 dilution. Incubate for 1 hour at room temperature.

Lot # _____ Exp. Date _____

For negative control slides, dilute mouse IgG1 control serum so that it's IgG1 protein concentration matches that of the primary antibody (if necessary). Then make a 1:25 dilution. If the concentrations can't be matched using this method, the dilution for the negative reagent may need to be adjusted.

Apply the negative and incubate for 1 hour at room temperature.

Lot # _____ Exp. Date _____

9. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

10. Apply the horse anti-mouse secondary antibody at a 1:500 dilution. Incubate for 30 minutes at room temperature.

Lot # _____ Date Reconstituted _____

11. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each.

12. Apply the Vectastain R.T.U. Elite Label and incubate for 30 minutes at room temperature.

Exp. Date _____ New Kit: yes / no

13. Rinse the slides in 2 changes of 1X wash buffer for 5 minutes each time.

14. Apply the DAB chromogen. Incubate in the dark for 6 minutes at room temperature.

(Add 1 drop of DAB per ml of substrate)

Lot # _____ Exp. Date _____ New Kit: yes / no

15. Rinse the slides in tap water 3 minutes.

16. Counterstain with hematoxylin for 20 seconds.

17. Rinse the slides in tap water until water is clear.

18. Gently agitate slides in 1X wash buffer until the tissues turn blue.

19. Dehydrate through the following solutions:

Solutions	Repetitions	Time
95% Ethanol	1 time	3 minutes
100% Ethanol	3 times	3 minutes
Xylene	2 times	5 minutes

20. Coverslip

Updated 03/17/11